


## Meteor Activity Outlook for February 22-28, 2025




Aaron Witbeck captured this brilliant fireball while photographing an intense aurora display on October 7, 2024, at 22:43 PDT (5:43 UT on Oct 8) from Yale, Washington USA. ©Aaron Witbeck

During this period, the moon reaches its new phase on Friday February 28th. On that date the moon will be located near the sun and will be invisible at night. This weekend the waning crescent moon will rise during the early morning hours but should not interfere with meteor observing as long as you keep the moon out of your field of view. The estimated total hourly rates for evening observers this weekend should be near 3 as seen from mid-northern latitudes (45N) and 4 as seen from tropical southern locations (25S). For morning observers, the estimated total hourly rates should be near 7 as seen from mid-northern latitudes (45N) and 12 as seen from tropical southern locations (25S). The actual rates seen will also depend on factors such as personal light and motion perception, local weather conditions, alertness, and experience in watching meteor activity. Morning rates are slightly reduced due to moonlight. Note that the hourly rates listed below are estimates as viewed from dark sky sites away from urban light sources. Observers viewing from urban areas will see less activity as only the brighter meteors will be visible from such locations.

The radiant (the area of the sky where meteors appear to shoot from) positions and rates listed below are exact for Saturday night/Sunday morning February 22/23. These positions do not change greatly day to day so the listed coordinates may be used during this entire period. Most star atlases (available online and at bookstores and planetariums) will provide maps with grid lines of the celestial coordinates so that you may find out exactly where these positions are located in the sky. I have also included charts of the sky that display the radiant positions for evening, midnight, and morning. The center of each chart is the sky directly overhead at the appropriate hour. These charts are oriented for facing south but can be used for any direction by rotating the charts to the desired direction. A planisphere or computer planetarium program is also useful in showing the sky at any time of night on any date of the year. Activity from each radiant is best seen when it is positioned highest in the sky, either due north or south along the meridian, depending on your latitude. Radiants that rise after midnight will not reach their highest point in the sky until daylight. For these radiants, it is best to view them during the last few hours before dawn. It must be remembered that meteor activity is rarely seen at its radiant position. Rather they shoot outwards from the radiant, so it is best to center your field of view so that the radiant lies toward the edge and not the center. Viewing there will allow you to easily trace the path of each meteor back to the radiant (if it is a shower member) or in another direction if it is sporadic. Meteor activity is not seen from radiants that are located far below the horizon. The positions below are listed in a west to east manner in order of right ascension (celestial longitude). The positions listed first are located further west therefore are accessible earlier in the night while those listed further down the list rise later in the night.



**Radiant Positions at 20:00 Local Standard Time**



**Radiant Positions at Midnight Local Standard Time**



## **These sources of meteoric activity are expected to be active this week**

The center of the large **Anthelion (ANT)** radiant is currently located at 11:08 (167) +06. This position lies in southeastern Leo, 3 degrees west of the 4th magnitude star known as sigma Leonis. Due to the large size of this radiant, Anthelion activity may also appear from western Virgo and northeastern Sextans as well as Leo. This radiant is best placed near 0100 local standard time (LST), when it lies on the meridian and is located highest in the southern sky. Rates at this time should be near 2 per hour no matter your location. With an entry velocity of 30 km/sec., the average Anthelion meteor would be of slow velocity.

The **26 Bootids (TSB)** were discovered by Zeljko Andreic and the Croatian Meteor Network team in 2014. These meteors are active from February 25 through March 8 with maximum activity occurring on March 4<sup>th</sup>. The radiant position lies at 14:05 (211) +07 on February 25<sup>th</sup>. This position lies on the Virgo/Bootes border, 6 degrees north of the 4<sup>th</sup> magnitude star known as tau Virginis. These meteors are best seen near 04:00 LST when the radiant lies highest in the southern sky. Hourly rates are expected to be less than 1 no matter your location. With an entry velocity of 49 km/sec., these meteors would possess a medium-fast velocity.

The **delta Normids (DNO)** were discovered by cameras of the All-Sky Meteor Surveillance System (CAMS) in their 2018 study of new radiants. This shower is active from February 13-March 17 with a peak on February 22. The radiant is currently located at 15:56 (239) -46, which places it in northwestern Norma, 2 degrees southwest of the faint star known as delta Normae. These meteors are best seen during the last hour prior to dawn when it lies highest above the southern horizon. Rates are expected to be less than 1 per hour no matter your location. With an entry velocity of 67 km/sec., these meteors would have a swift velocity. Due to the southern location of this radiant, these meteors are not well seen north of the northern tropics. These meteors may be synonymous with the gamma Normids, which were thought to be active in mid-March.

**Sporadic** meteors are those meteors that cannot be associated with any known meteor shower. All meteor showers are evolving and disperse over time to the point where they are no longer recognizable. Away from the peaks of the major annual showers, these sporadic meteors make up the bulk of the activity seen each night. As seen from the mid-northern hemisphere (45N) one would expect to see during this period approximately 5 sporadic meteors per hour during the last hour before dawn as seen from rural observing sites. Evening rates would be near 2 per hour. As seen from the tropical southern latitudes (25S), morning rates would be near 10 per hour as seen from rural observing sites and 3 per hour during the evening hours. Locations between these two extremes would see activity between these listed figures. Morning rates are reduced due to moonlight.

The list below offers the information in tabular form of the showers that I feel are within reach of the visual observer to discern. Hourly rates are often less than one, so these sources are rarely listed as visual targets in most meteor shower lists. If you are like me and wish to associate as many meteors as possible with known sources, then you will appreciate these listings. Before listing meteors from these obscure sources, you should attempt to prove these meteors belong to them and are not chance alignments of sporadic meteors. You can note parameters such as duration, length, radiant distance and the elevation of each meteor to help compute the probability of shower

association. It should be remembered that slow meteors can be seen from fast showers, but fast meteors cannot be produced from slow showers. Slower showers are those with velocities less than 35/km per second. Slow meteors can appear from fast showers when they appear close to the radiant or low in the sky. The table located on page 22 of the [IMO's 2025 Meteor Shower Calendar](#) is a big help in aiding in the identification of meteors. If you record the length and duration of each meteor, you can use this chart to check the probability of the meteor belonging to a shower of known velocity. If the angular velocity is similar to the figure in the table, then your meteor probably belongs to that shower. Rates and positions are exact for Saturday night/Sunday morning.

SHOWER	DATE OF MAXIMUM ACTIVITY	CELESTIAL POSITION	ENTRY VELOCITY	CULMINATION	HOURLY RATE	CLASS
		RA (RA in Deg.) DEC	Km/Sec	Local Standard Time	North-South	
Anthelion (ANT)	-	11:08 (167) +06	30	01:00	2 - 2	II
26 Bootids (TSB)	Mar 04	14:05 (211) +07	49	05:00	<1 - <1	IV
delta Normids (DNO)	Feb 22	15:56 (239) -46	67	07:00	<1 - <1	IV

You can keep track of the activity of these meteor showers as well as those beyond the limits of visual observing by visiting the [NASA Meteor Shower Portal](#). You can move the sky globe to see different areas of the sky. Colored dots indicate shower meteors while white dots indicate sporadic (random) activity. The large orange disk indicates the position of the sun so little activity will be seen in that area of the sky.

**Class Explanation:** A scale to group meteor showers by their intensity:

- **Class I:** the strongest annual showers with Zenith Hourly Rates normally ten or better.
- **Class II:** reliable minor showers with ZHR's normally two to ten.
- **Class III:** showers that do not provide annual activity. These showers are rarely active yet have the potential to produce a major display on occasion.
- **Class IV:** weak minor showers with ZHR's rarely exceeding two. The study of these showers is best left to experienced observers who use plotting and angular velocity estimates to determine shower association. These weak showers are also good targets for video and photographic work. Observers with less experience are urged to limit their shower associations to showers with a rating of I to III.