


# Journal of the Association of Lunar & Planetary Observers



*The Strolling Astronomer*

Volume 43, Number 3, Summer 2001

Now in Portable Document Format (PDF) for MacIntosh and  
PC-Compatible Computers

*Inside . . .*


## *Our VERY active Sun*

It's been a most rewarding summer for solar observers, whether they use h-alpha or solar screen or aluminized mylar filters.

This month's cover features a dramatic astrophoto of a powerful (X-class) solar flare, taken 25 October, 2001, 15:46 UT by Frank Schiralli, Jr. of Northport, NY. Contact him at

[astroimager11768@yahoo.com](mailto:astroimager11768@yahoo.com)

**Equipment: Coronado 40mm SolarMax filter, 94mm Brandon, and Starlight Xpress MX916 camera.**



**Also** ..a follow-up report on hook-like shadows on the floor of the lunar crater Plato, reports on Venus, the Remote Planets, and an index to Volume 42 of the Strolling Astronomer -- plus ALPO observing section reports and much, much more.

# Journal of the Association of Lunar & Planetary Observers, The Strolling Astronomer

Volume 43, No. 3, Summer 2001

This issue published in October 2001 for distribution in both portable document format (pdf) and also hardcopy format.

This publication is the official journal of the Association of Lunar and Planetary Observers (ALPO).

The purpose of this journal is to share observation reports, opinions, and other news from ALPO members with other members and the professional astronomical community.

© 2001, Association of Lunar and Planetary Observers (ALPO). The ALPO hereby grants permission to educators, academic libraries and the professional astronomical community to photocopy material for educational or research purposes as required. There is no charge for these uses provided that credit is given to *The Strolling Astronomer*, the "JALPO" or the ALPO itself, as appropriate. All others must request permission from the ALPO.


For membership or general information about the ALPO, contact:

Matthew Will  
ALPO Membership Secretary/Treasurer  
P.O. Box 13456  
Springfield, Illinois 62791-3456

E-mail to: will008@attglobal.net

Visit the ALPO online at:

<http://www.lpl.arizona.edu/alpo>



## In This Issue:

Point of View ..... 1

## The ALPO Pages

Letters .....	2
Solar Section Report.....	2
Mercury Section Report .....	3
Venus Section Report .....	4
Lunar Section Report	
Selected Areas Program .....	4
Lunar Meteoritic Impact Search.....	5
Meteors Section Report .....	6
Mars Section Report .....	6
Jupiter Section Report .....	7
Saturn Section Report.....	7
Remote Planets Report .....	7
ALPO Website Report .....	8
Other Contributions .....	8
About the Authors .....	9
Errata .....	10

## Features

ALPO Solar Section, Solar Activity Report: Carrington Rotations 1911-1922.....	11
ALPO Observations of Venus: The 1998 Western (Morning) Apparition .....	17
The Nature of the Hook-Like Shadow on Plato's Floor .....	24
Observations of the Remote Planets in 2000 .....	30
Index to Volume 42 of <i>The Strolling Astronomer</i> .....	38

## ALPO Resources

ALPO Board of Directors.....	41
Publications Staff.....	41
Lunar and Planetary Training Program .....	41
Observing and Interest Section Staff ..	41-43
ALPO Board/Staff E-mail Directory .....	43
ALPO Publications:	
The Monograph Series .....	43
ALPO Observing Section Publications.	44
Other ALPO Publications .....	45

## **Association of Lunar and Planetary Observers (ALPO)**

### **Board of Directors**

Executive Director (Chair); Julius L. Benton, Jr.  
Associate Director, Donald C. Parker  
Founder/Director Emeritus; Walter H. Haas  
Member of the Board; Richard Hill  
Member of the Board; Ken Poshedly  
Member of the Board; Richard W. Schmude, Jr.  
Member of the Board; John E. Westfall  
Member of the Board, Secretary; Elizabeth W. Westfall  
Member of the Board, Treasurer and  
Membership Secretary; Matthew Will

### **Publications**

Acting Publisher, Ken Poshedly

### **Primary Observing Coordinators, Other Staff**

(See full listing in *ALPO Resources* at end of book)

#### **Lunar and Planetary Training Program Coordinators:**

Timothy L. Robertson and Matthew Will

**Solar Section:** Coordinator, *Website, SolNet, Rotation Report*, handbook; Richard Hill

**Mercury Section:** Acting Coordinator; Frank Melillo

**Venus Section:** Coordinator; Julius L. Benton, Jr.

**Mercury/Venus Transit Section:** Coordinator; John E. Westfall

**Lunar Section:** Coordinator; *Selected Areas Program*; Julius L. Benton, Jr.

**Mars Section:** Coordinator, *all observations, U.S. correspondence*; Daniel M. Troiani

**Minor Planets Section:** Coordinator; Frederick Pilcher

**Jupiter Section:** Acting Coordinator; Richard W. Schmude, Jr.

**Saturn Section:** Coordinator; Julius L. Benton, Jr.

**Remote Planets Section:** Coordinator; Richard W. Schmude, Jr.

**Comets Section:** Coordinator; Gary Kronk

**Meteors Section:** Coordinator; Robert D. Lunsford

**Meteorites Section (provisional):** Acting Coordinator; Dolores Hill

**Computing Section:** Coordinator; Mike W. McClure

**Youth Section:** Coordinator; Matthew Will

**Historical Section (provisional):** Acting Coordinator; Richard Baum

**Instruments Section:** Acting Coordinator; R.B. Minton

**Eclipse Section (provisional):** Acting Coordinator; Michael D. Reynolds

**Webmaster:** Coordinator; Richard Hill

### **Point of View**

**By Ken Poshedly, Acting Editor & Publisher  
Journal of the Assn. of  
Lunar & Planetary Observers**



Even though I may be criticized for saying so, I think I would be remiss in not recognizing that the terrorist attacks of Sept. 11 touched nearly every facet of our lives, including those of us in the Assn. of Lunar & Planetary Observers.

Whether you live in California or New York, or even overseas, and even if you knew no one at the World Trade Center, the Pentagon or on the planes that carried the hundreds of people to their deaths, the ripple effects of the terrorist attacks continue to trickle down and affect just about all we do. Today, all of us are much more wary about the envelopes we get in the mail, the people we see in lines next to us in public places, and we may even be a little more careful about our jaunts to dark sky sites for sky observing and ALPO sessions.

While I myself knew no one firsthand who was lost or injured, a fellow member of the Atlanta Astronomy Club here lost his nephew, Jeremy Glick, who was one of those brave persons who used his cellphone to let the world know his plane had been hijacked and that the passengers were at least going to try to take it back. We'll never know for sure, but maybe it was the actions of Jeremy and his fellow passengers that kept the plane from reaching Washington, DC. Instead, it crashed outside Pittsburgh, Pennsylvania.

And our good members who live in or anywhere near New York City continue to suffer the direct pain of that tragic day as the site of the former office building complex is cleared, bodies are found, and even smoke and odors from the ongoing fires continue to waft far away.

One wonders how can they even think of hoisting a telescope out anymore. But to some, our interests and our celestial activities have therapeutic value. Some folks use astronomy partly as a means to clear their minds of sad and tragic events. They don't get the satisfaction that once was. But they know that it will eventually return. Right now, astronomy may be the only thing that keeps them from falling apart themselves.

We, the ALPO, offer our deepest sympathies and condolences to those of you who lost loved ones in the Sept. 11 tragedies.

## The ALPO Pages: Member, section and activity news

### Letters

#### Reference follow-up needed

... on page 29 ("Observing for Lunar Flashes"), there is a reference to an article by Stuart Leon as being in the JALPO Vol. 10, Nos. 5-6, pp. 42-43, 1956. I cannot find the article in my copy. Could you please have the author research this and publish a correction in the next issue? In the mean time, please let me know what is the correct reference. BTW. I now have a near complete run of the JALPO back to the beginning. I am missing only a few issues.

-- Bob Garfinkle, [ragarf@earthlink.net](mailto:ragarf@earthlink.net)

#### Equation clarification

In his very valuable article on the extensions of the cusps of Venus (JALPO 42(2):58-66, April, 2000), Mr. Walter Haas calculates theoretical extensions of the cusps using the equation,

$$\cos(p/2) = -\sin l \csc i \quad (\text{his equation 3})$$

He assumes that this equation is inexact because it is an approximation of his unsolvable equation 1. He then proposes (p. 63, list item 4) that one reason the theoretical cusp extensions do not correspond precisely to observations is that, due to the approximate nature of equation 3, the calculations are merely approximations.

In fact, Haas's equation 3 is precise. I have developed a four-page proof of this that I am happy to send to anyone who sends a self-addressed, stamped envelope to me at 3405 Woodstone Place, Augusta, GA 30909. The proof is of too narrow interest to merit publication in JALPO.

The thrust of Haas's article is unaffected by this proof. The only one of his conclusions that should be changed is his supposition that more accurate theoretical calculations might correspond more closely to the observed values.

Roger Venable, [rjvmd@knology.net](mailto:rjvmd@knology.net)

Look for details of the ALPO Board meeting in your next *Strolling Astronomer*

### Thinking of contributing?

The ALPO welcomes reports, articles, and letters for publication. In most cases, such materials are considered for the Journal of the ALPO (*The Strolling Astronomer*); the Guidelines that follow are intended for that Journal. There are also special guidelines for ALPO monographs and proceedings.

- All submissions, including "Letters to the Editor", should be accompanied by a cover letter stating that the material is being submitted for publication. Please give your telephone number, fax number and e-mail address if you have them. A single copy of a submission is usually sufficient, although we recommend strongly that the author(s) retain their own copy.
- In terms of content, we deal with the objects in the Solar System; the Sun, Moon, major and minor planets, comets, and meteors. Our emphasis is on observations and observing techniques rather than theory or cosmology. Ordinarily, submissions should not exceed a total of 10-12 printed pages including illustrations; allow about 1,000 words per printed page.
- The ALPO publishes in the English language only. All submissions should be in clear, grammatical English.

The full set of publication guidelines can be found online at <http://www.lpl.arizona.edu/~rhill/alpo/pubguide.html>

### Solar Section

#### By Rik Hill

THE LEONIDS ARE COMING! In only a couple weeks we will all be waiting breathless for the possibility of a meteor storm. The latest predictions for this activity will be posted at:

<http://www.lpl.arizona.edu/~rhill/alpo/meteorstuff/outlook.html>

The Lunar Meteoritic Impact Search, a sub-Section of the ALPO Lunar Section now in it's second year, is mounting a big effort to observe some Leonid impacts on the moon. Get the details at:

<http://www.lpl.arizona.edu/~rhill/alpo/lunarstuff/lunimpacts.html>

Jupiter observations continue to pour in as the grand planet approaches opposition. Incredible work is now routinely being done by amateur astronomers at or near the resolution limits of their instruments. We invite you to take a look at some of the results at:

<http://www.lpl.arizona.edu/~rhill/alpo/jupstuff/recobs.html>

Though Mars is starting to get small, still there are a number of amateurs that are still imaging the planet. Their work can be enjoyed at:

<http://www.lpl.arizona.edu/~rhill/alpo/marstuff/recobs.html>

The Remote Planets Section, Recent Observations page has had a lot of upgrading and additions. Here you will find some spectacular work done with modest apertures in various wavelengths and spectroscopy!

<http://www.lpl.arizona.edu/~rhill/alpo/remplan.html>

There's still a lot of solar activity to be had and you can see some of it on the Solar Section pages at:

<http://www.lpl.arizona.edu/~rhill/alpo/solar.html>

Go to the Recent ALPOSS Observations link for up to date observations by amateur astronomers from around the world.

## Mercury Section

### By Frank Melillo

(The following are excerpts from his complete report which can be found on the ALPO website.)

#### COORDINATOR'S CORNER:

The ALCON/ALPO convention took place in Frederick, MD, on July 25 - 28, 2001, and it was very well-attended. It was there that I presented my talk on "Scientific Interests of the Planet Mercury". (Lots of compliments afterwards on my images of Mercury.)

It wasn't until after I left (early, for family reasons) that I learned that I had won the ALPO "Walter Haas" award in recognition of many years of excellent planetary observations.

I met many ALPO member at the convention. It was such a thrill that I have corresponded with them for many years and finally I met them in person. In fact, I heard that it was the first time that the ALPO convention papers covered Mercury through Pluto. Both planets are rarely mentioned in many talks. As you can see, Mars,

Jupiter and Saturn are very well-observed. Most importantly, the remote planets and even Mercury and Venus now need more attention.

#### MERCURY NEWS:

As stated in my last newsletter, the morning apparition (June 16 to August 6) was a so-so opportunity to observe Mercury. But I did take some excellent images of Mercury. The best so far! Even Mario Frassati of Italy, did some excellent sketches of Mercury. In fact, Mario and I were the only ones to contribute during that apparition.

On July 14, 15 and 21, I took images of Mercury in full daylight about 70 degrees above the horizon! The observations were done when the sky was crystal blue.

Thanks to Venus! I used it as a guide to find Mercury. After I set up Venus' coordinates, it took me 30 seconds to find Mercury in a low-power field. Believe me, it wasn't too difficult at all! Mercury's phase was clear as a bell. I replaced the eyepiece with a CCD camera and took some excellent shots.

Johan Warell of Uppsala University in Sweden made some excellent researches on Mercury for the past 5 years. He even wrote a paper on "Properties of the Herman regolith: Global regolith albedo variation at 200km scale from multicolor CCD imaging". I tell you that this is one excellent paper! At one point, he made a synthesized map of Mercury on July 14. With his albedo map on that date, Mercury should appear like this:



Synthesized Mercury disk map (right) by John Warell with comparison of ccd image (left) by Frank Melillo. Taken 14 July, 2000, CM = 263°

Believe it or not, I had imaged Mercury at the very same day! What I did was I to degrade the map's resolution, making it equivilant in scale to an image as seen in an 8-inch telescope. Surprisingly, both images -- his and mine -- look identical to each other!

## Venus Section

### By Julius Benton

ALPO Venus Section observers attempt to follow Venus on every possible clear night throughout an apparition of the planet, which runs from conjunction to subsequent conjunction with the Sun.

The minimum recommended aperture for useful observations of Venus is about 15.2 cm. (6.0 in.) for reflectors or 7.5 cm. (3.0 in.) for refractors. Low-transmission color filters increase contrast and definition while limiting the effects of irradiation. For example, Wratten blue (W38A) and violet (W47) filters are useful for low contrast detail, mainly because of the yellowish atmospheric cloud layer. Variable-density polarizing filters are especially useful in reducing glare and improving the visibility of faint markings.

Since Venus is an inferior planet (with an interior orbit to that of the Earth), it exhibits phases like the Moon. Also, since Venus is comparatively near the Sun, it is characteristically very bright. The high albedo produces an excessive amount of glare, making it difficult for observers to see elusive, low-contrast atmospheric markings on Venus. Controversy persists as to the true nature of these dusky amorphous or somewhat streaky atmospheric features. It is not at all unusual for two observers, working on the same date with comparable instrumentation, to see striking dissimilar atmospheric phenomena on the planet.

Venus attains a maximum angular distance of about 47° from the Sun, so it is usually observed near sunrise or sunset. When Venus is seen against a dark sky, the effects arising from its excessive brilliance are pronounced. At eastern (evening) apparitions, Venus is typically low in the western sky where atmospheric refraction and prismatic dispersion can ruin image quality, and many observers try to view Venus only when it has an altitude of 20° or more above the horizon. At times of western (morning) apparitions, observers can wait until Venus gains altitude and the background sky brightens considerably. Also at such times, Venus can readily be followed into daylight when most of the prevailing glare associated with the planet is reduced.

The ALPO Venus Section international observing network is currently comprised of about 20-25 observers, and there is active professional-amateur collaboration. Widely-spaced observations are of limited value, and the ALPO Venus Section stresses systematic, as well as simultaneous, observations to improve the objectivity of reports of elusive atmospheric features on Venus. Drawing blanks and report forms are furnished by the ALPO Venus Section to record data, and studies of Venus are organized into the following main programs:

- Visual observation and categorization of atmospheric details in dark, twilight, and daylight skies.

- Drawings of atmospheric phenomena.
- Observation of cusps, cusp-caps, and cusp-bands, including defining the morphology and degree of extension of cusps.
- Observation of dark hemisphere phenomena, including monitoring the visibility of the Ashen Light.
- Observation of terminator geometry (monitoring any irregularities).
- Studies of Schroter's phase phenomenon.
- Visual photometry and colorimetry of atmospheric features and phenomena.
- Routine photography (UV photography is stressed), CCD imaging, photoelectric photometry, and videography of Venus.

Individuals interested in participating in the ALPO Venus programs should contact:

Julius L. Benton, Jr., Ph.D.  
Coordinator, ALPO Venus Section  
Associates in Astronomy  
305 Surrey Road  
Savannah, GA 31410

E-Mail: [jlbaina@msn.com](mailto:jlbaina@msn.com) (general correspondence)

E-Mail: [jbenton55@home.com](mailto:jbenton55@home.com) (image files and attachments)

## Lunar Section: Selected Areas Program

### By Julius Benton

The ALPO Lunar Selected Areas Program stresses systematic visual monitoring of variable phenomena at the surface of the Moon with specific emphasis on suspected long-term cyclical or "seasonal" albedo or morphological fluctuations in "*selected areas*" over many lunations. This is one of many areas of lunar research in which the skillful and imaginative amateur astronomer can find worthwhile observational opportunities.

Special talents for drawing lunar features, although definitely helpful, are not necessary for this program. The most fundamental and essential prerequisite for participation in the Selected Areas Program is the willingness to follow the Moon and the chosen feature(s) for many consecutive lunations, year after year. A good 10.2cm. (4.0in.) refractor or 20.0cm. (8.0 in.) reflector will deliver sufficient resolution of lunar detail for full participation in nearly all aspects of the program.

Initially, the observing program involves monitoring selected lunar features to establish a baseline of the *normal*, yet often complex, changes in appearance and albedo that they undergo from lunar sunrise to sunset over many lunations. This "control" serves as a reference for observers so they might recognize *anomalous* variations in morphology, tone (albedo), or color, that

are attributable to changing solar angle (phase angle) or libration, and which clearly *do not* repeat systematically from lunation to lunation. Anomalous phenomena that the Selected Areas Program looks for are:

1. *Tonal and/or Color Variations.* These are variations in tone or color, or in the size and shape of a region of tone or color, that is not related to changing illumination (i.e., the phenomenon does not exactly repeat from lunation to lunation). Areas in lunar features most subject to such anomalous behavior are *radial bands, dark patches, and nimbi or haloes.*

2. *Shape and Size Changes.* These are variations in the appearance and morphology of a feature that cannot be traced to changing solar illumination or libration.

3. *Shadow Anomalies.* These are deviations of lunar shadows away from the theoretical normal absolute black condition, or a shadow with an anomalous shape or hue, in most cases not attributable to changing phase angle.

4. *Appearance or Disappearance of Features.* Although exceedingly improbable and controversial, these are features that seem to be present now, but appear to be absent on earlier maps or photographs; or, features that are no longer visible today but which are clearly indicated on earlier maps or photographs.

5. *Features Exposed to Earthshine.* These are any anomalous tonal or albedo phenomena (any of the categories listed here) that occur under the conditions of Earthshine.

6. *Eclipse-Induced Phenomena.* These are features that exhibit anomalous characteristics (categories 1 through 4 above) during and after an eclipse, compared with previous eclipses when the same areas were monitored.

The lunar features that are currently designated as the *official* lunar formations that are being monitored as part of the Selected Areas Program (SAP) are *Alphonsus, Aristarchus-Herodotus, Atlas, Copernicus, Plato, Theophilus,* and *Tycho.* All of these areas were chosen because they are relatively easy to find, convenient to observe, and have historically shown numerous instances of suspected anomalies. Complete outline charts and observing forms are available from the ALPO Lunar Section for each of the features noted.

Thus, observations of the Moon that are specific to the Lunar Selected Areas Program may be summarized as:

- Visual photometry of specific lunar features, defining their normal albedo profiles throughout a lunation as a function of changing solar illumination.
- Visual photometry of specific lunar features, monitoring variations from their normal albedo that are not simply a result of changing solar illumination.

- Drawings of specific lunar features throughout a lunation and from lunation-to-lunation in conjunction with visual photometry.
- Routine photography, CCD imaging, photoelectric photometry, and videography of specific lunar features to supplement visual photometry programs throughout a lunation and from lunation-to-lunation.
- Comparative analysis of lunar features and albedo profiles.

It should be pointed out that the Selected Areas Program now includes the Bright and Banded Craters Program and the Dark-Haloed Craters Program. Further information on these programs is available upon request. Complete details on our observing programs can be found in the *ALPO Lunar Selected Areas Handbook.* Individuals interested in participating in the ALPO Lunar Selected Areas Program should contact:

Julius L. Benton, Jr., Ph.D.  
Coordinator, ALPO Lunar Section  
Selected Areas Program  
Associates in Astronomy  
305 Surrey Road  
Savannah, GA 31410

E-Mail: [jlbaina@msn.com](mailto:jlbaina@msn.com) (general correspondence)

E-Mail: [jbenton55@home.com](mailto:jbenton55@home.com) (image files and attachments)

## **Lunar Section: Lunar Meteoritic Impact Search Report**

**By Brian Cudnik**

Regular monitoring of the Moon continued during 2001, but focused more on the the flux of meteoroids received by the waxing crescent Moon during a monthly campaign carried out by two individuals. Several other individuals contributed observations of possible impact candidates, but none of them has been confirmed to date. Our focus shifts toward the Leonid event in November, which could possibly be the last significant opportunity to observe lunar meteorites for quite some time. There is always opportunity to watch for lunar meteorites, but the probability of observing and confirming them is greatest during the Leonids due to the combination of high meteoroid flux and velocity.

Dr. Anthony Cook of Washington, D.C., in January began a systematic watch of the Moon for a period of one week each month. This "Earthshine watch" occurred monthly during the period from three days after New Moon to two days after First Quarter. He was joined in the effort by Sam Falvo of Utica, New York, in May. Several candidate events were reported by these two,

and two additional observers reported events as well. The vast majority of Dr. Cook's many observed "events" were actually cosmic ray signatures. His candidates which have the greatest likelihood of being genuine events appear below. However, it is important to stress that none of the events listed has been independently confirmed, so it is very possible that each of these is artificial as well.

Some of Tony Cook's best candidates:

- April 30th, 01:55:45UTC "I'd go for as a lunar impact on this night - this one was merged with background noise and about the size of Aristarchus."
- Apr 27, 00:43:25 UTC - "[I]t is difficult to be tell if this a 'cosmic ray or impact flash' for sure"
- Apr 27 01:12:02 UTC - "[I]t is difficult to tell if this a 'cosmic ray or impact flash' for sure?"
- 01:18:29, 27 Aug UTC - "Bright flash S of Timocharis. Size and brightness not too disimilar to nearby sunlit peaks on terminator.

Sam Falvo's candidate:

02:55:03, 28 Jun 28 UT - "a brief flash, much less than one-second duration. It was in the dark side to the left of partially illuminated Hipparchus. I estimate it to be quite near the crater Flammerion."

Other impact candidates were observed by IOTA member Dave English, of Oceanside, California, at 6:58:30 UT, 30 May 30; and Capt. R. Patterson (ret) of Port St John, Florida, on 26 June 2001, 3:39 UT

Dr. Cook has come to an initial conclusion that during non-shower times, the rate of capture of genuine impact events on the Moon is about one per tens of hours of observing time. Further observations by him and other LMIS observers will help to improve this assertion and obtain a more refined value.

## **Meteors Section**

### **By Robert Lunsford**

The ALPO Meteors Section continues to provide current information on meteor activity through two sources: the Meteors Section Newsletter (published quarterly) provides a review of the meteor activity seen during the most recent quarter with articles and actual data from observers. It also provides an outlook for the next quarter. The Meteors Section website provides an in depth analysis of the upcoming week's meteor activity plus a general outlook for the current year.

The section coordinator is participating in an expedition to Arizona to observe and record the expected Leonid storm this November. The following month, he and the assistant coordinator will lead another expedition to the Mojave Desert to view the Geminid meteor shower. Reports on these expeditions will be available in the

December 01 and the March 02 issues of the ALPO Meteors Section Newsletter.

## **Mars Section**

### **By Daniel P. Joyce, Assistant Coordinator, Mars Section Dust Subsiding???**

The dust activity that has been the dominant phenomenon witnessed by hundreds of observers since the fourth week of June has at least shown signs of thinning out in recent weeks. It may yet be too early to draw the conclusion that by the end of October all albedo features will yet again attain their normal prominence. Even as some regions have slowly been unveiled, there appear to be possible new dust outbreaks as the favorable season has been reached. It is once again important to assert that the June outbreak was entirely uncharacteristic of the season. This demonstrates the alertness and discernment on the part of those who (mostly in Japan) did not allow preconceived notions to impair their perception of what they were seeing and of having the shrewdness to immediately report their findings.

In the not-too-distant-future, it is hoped, the Section will begin to acknowledge the fine work of the observing network that spans the globe. It was a particularly difficult time for our European friends, especially north of Italy, as often a 10 degree maximum altitude for Mars had to suffice. We are indebted to those who provided imagery despite the conditions! Probably not enough could ever be said about the specular reflection sightings made by a party of observers who not only saw them, but were the beneficiaries of a prediction to the effect that the phenomenon would in fact occur. Research into past observations, even those scant (and arguably obscure) paid huge dividends. The observations of nearly a half-century ago were painstakingly correlated with modern space probe accumulated data when it was not obvious that such a connection was there for the making.

It is also to be noted that a German periodical "Sterne Und Weltraum" (Stars and World?) has placed Mars prominently on its July cover and provided a substantial accompanying pull-out chart which they attribute in some measure to the work of our Section coordinator Dan Troiani (Mars Global Surveyor helped a little, too!).

But don't quit just yet; the apparition is far from over and Mars is still ten arc-seconds. Furthermore, it is just past perihelion and therefore able to maintain a fair pace with the faster earth, which therefore cannot pull away at its customary rate at this date past opposition. So keep looking!

## Jupiter Section

### By Richard Schmude

The 2000-2001 Jupiter report is almost complete. The magnitudes and color indexes of Jupiter were measured by R. Schmude and H. Lesser in late 2000, and the resulting magnitudes were close to the expected value. Dr. Schmude also measured the brightness of Jupiter over a 5-hour period on Dec. 8 and found that Jupiter became about 2 - 3% dimmer as the Great Red Spot crossed the central meridian. In mid-2001, Jupiter became visible in the morning sky. The biggest changes are that the North Equatorial Belt became dimmer and that the Equatorial Zone became more yellow.

## Saturn Section

### By Julius Benton

Almost any optical assistance will show Saturn's spectacular ring system, and the major disc features are revealed with a 7.5cm. (3.0 in.) refractor, including perhaps a major belt and a zone or two near the equator of the planet. Cassini's Division should also be visible in the rings with such an instrument. With a 10.2cm. (4.0 in.) refractor or a 15.2cm. (6.0 in.) reflector, the observer will discover that he has found about the minimum aperture suitable for routine and beginning detailed studies of Saturn. Of course, when seeing conditions allow, the larger the aperture, the greater will be the image scale and the better the resolution and image brightness.

A novice should spend at least some time in experimental work with the telescope he intends to use for routine views of Saturn. One should always try to establish the best combination of magnification, filters, and image size, brightness, and contrast. These topics, and many others, are discussed in considerable detail in *The Saturn Handbook*. After gaining some practical experience in observing Saturn, individuals are encouraged to become familiar with more advanced methods and techniques for observing.

Like Jupiter, Saturn displays in an appropriate telescope a series of bright zones and darker belts that run roughly parallel to the equator. Much of the fundamental nomenclature assigned to the specific zones and belts of Jupiter applies to Saturn. *The Saturn Handbook* gives more detail on the specialized uses of all of the terminology and nomenclature for Saturn's global and ring features.

The ALPO Saturn Section international observing network is currently comprised of about 30-35 observers, and there is active professional-amateur collaboration. Studies of Saturn's globe, rings, and satellites are organized into the following main programs:

- Visual numerical relative intensity estimates of belts, zones, and ring components.

- Full-disc drawings and sectional sketches of global and ring phenomena (the Saturn Section furnishes templates with the correct global oblateness and ring geometry to facilitate making drawings).
- Central meridian (CM) transit timings of details in belts and zones on the globe of Saturn (utilized to determine or confirm rotation rates in various latitudes).
- Visual latitude estimates or filar micrometer measurements of belts and zones on the globe of Saturn.
- Colorimetry and absolute color estimates of globe and ring features.
- Observation of "intensity minima" in the rings (in addition to observations of Cassini's and Encke's divisions).
- Observational monitoring of the bicolored aspect of the rings of Saturn.
- Observations of stellar occultations by Saturn's rings.
- Specialized observations of Saturn during edgewise ring presentations in addition to routine studies.
- Visual observations and magnitude estimates, as well as photoelectric photometry, of the satellites of Saturn.
- Spectroscopy and photometry of Titan.
- Routine photography, CCD imaging, photoelectric photometry, and videography of Saturn and its ring system.
- Simultaneous observations of Saturn.

Individuals interested in participating in the ALPO Saturn programs should contact:

Julius L. Benton, Jr., Ph.D.  
Coordinator  
ALPO Saturn Section  
Associates in Astronomy  
305 Surrey Road  
Savannah, GA 31410

E-Mail: [jlbaina@msn.com](mailto:jlbaina@msn.com) (general correspondence)

E-Mail: [jbenton55@home.com](mailto:jbenton55@home.com) (image files and attachments)

## Remote Planets Section

### By Richard Schmude

About 10 people have submitted a variety of measurements of Uranus, Neptune and Pluto during the past 6 months. Doug West has measured the brightness of Uranus, Neptune and Pluto using a CCD camera along with a Johnson V-filter. Dr. Schmude has measured the brightness of Uranus and Neptune and the results indicate that Uranus continues to get slightly dimmer. Neptune's brightness is about the same as what it was last year. Frank Melillo has measured the relative brightness

of Titania, Oberon and Umbriel. Umbriel was measured to be 1.02 magnitudes dimmer than Oberon. Antonio Cidadao in Portugal was able to measure the Titania occultation on Sept. 8, 2001. R. Schmude and F. Melillo also carried out measurements but did not observe an occultation.

## ALPO Website News

### By Rik Hill

Mercury is now leaving the evening sky but you can still enjoy it with the latest *Mercury Today*, now available at:

<http://www.lpl.arizona.edu/~rhill/alpo/mercstuff/newsletter.html>

Frank Melillo, has also just posted his latest Mercury images and Mercury newsletter on the ALPO Mercury Section pages at:

<http://www.lpl.arizona.edu/~rhill/alpo/merc.html>

You cannot help but be impressed by his work with only an 8-inch aperture and why he is so deserving of the award he was given.

The Minor Planets Section's Magnitude Alert Project is active and making regular postings to:

<http://www.lpl.arizona.edu/~rhill/alpo/minplan/alert2001.html>

And the ALPO Meteors Section makes its regular announcements and alerts to observers at:

<http://www.lpl.arizona.edu/~rhill/alpo/meteorstuff/outlook.html>

Both of these pages keep ALPO observers up-to-date with the latest information in their respective studies.

The June solar eclipse has spawned a bunch of great images by ALPO observers. You can see some of these at:

<http://www.lpl.arizona.edu/~rhill/alpo/eclstuff/010621.html>

with more to come! And while you're there browse the rebuilt Eclipse Section page at:

<http://www.lpl.arizona.edu/~rhill/alpo/eclipse.html>

Mars is now at east quadrature. Though its diameter is only 11 arc sec. it is still very observable. The numbers of observations submitted to the ALPO exceeds that of the last apparitions and has caused a reworking of the Mars Section observations page at:

<http://www.lpl.arizona.edu/~rhill/alpo/marstuff/marsalert.html>

Since the reworking, this page loads more rapidly and the user has quicker access to images posted.

Thanks to the generosity of Sylvain Rondi (France) there's a new look to Solar Section webpage. He has given us permission to use one of his solar prominence animations on our page and it really does look good:

<http://www.lpl.arizona.edu/~rhill/alpo/solar.html>

The Sun is still extremely active, and you can keep up with the activity at the newly rebuilt Solar Section observations page at:

<http://www.lpl.arizona.edu/~rhill/alpo/solstuff/recobs.html>

Thanks to the work and generosity of Patrick McIntosh (*HelioSynoptics*) we also have graphic analysis of recent activity at the top of the ALPO solar webpage.

For a more comprehensive summary of ALPO activities, subscribe to our quarterly journal the *Strolling Astronomer - Journal of the ALPO* (JALPO). This is now available in digital format (DJALPO) at a reduced rate with faster delivery. See our website for details.


## Other Contributions

These contributions are presented here to demonstrate what others are doing and hopefully to inspire others to do the same. These reports and images have not undergone ALPO peer review for scientific study. All ALPO members are invited to send their contributions directly to the ALPO section coordinator for the subject at hand. Also, due to the highly subjective nature of color sketches and drawings, only black & white drawings can be presented. Color astro-images which more objectively represent the true nature of the subject will be presented here.

### Saturn

An observing report and sketch on Saturn from Eric Jamison <ericj@metro2000.net> of Pepperell, Massachusetts, posted on the *Shallow-Sky* listserv:

October 5, 2001, 8:25 - 9:05 UT, seeing 7 - 8, transparency 2. Astro-Physics 5.1" f/8.35 EDF refractor home-made Dobsonian-style mount. Magnification 264 - 308x with AP/Baader/Zeiss binoviewer.




The shadow of the globe was visible on the rings, giving

the planet a 3-dimensional look to it, but not as pronounced as it had been in early September. In the Equatorial Zone (EZ), the Equatorial Band (EB) was visible. Shading was noted on both the preceding and following ansae of the B-Ring. The C or Crepe Ring prominent, and the EZ north (EZn) was faintly visible through the C Ring. A-Ring appeared darker than the B-Ring.

The left (or preceding) ansa of the A-Ring showed some variation in tone, with lighter and darker areas visible. On the right (or following) ansa, the lighter areas were not visible and the darker areas were less pronounced. These darker areas were somewhat similar to an earlier observation I had of Saturn in September 1998, but smaller. Five satellites were visible near Saturn, including Tethys, Dione, Rhea, Titan, and Iapetus.

## Sun

From Ginger Mayfield of Divide, Colorado <chikadee@earthlink.net>, posted on the *SolNet* list-serv:



“The weekend at the Great Plains Star Party (16 Oct.) I had a chance to do some solar viewing through a new friend's 90mm refractor and Coronado H-alpha filter. It was one of those beautiful blue sky

days that comes after cloudiness and we were all up for some sunshine and solar viewing. After viewing with a single eyepiece for awhile we decided to try my TV binoviewer and the views were really amazing! There were several large prominences and the surface detail almost seemed three dimensional with the binoviewer. We were lucky to have great weather all afternoon and many folks came by for a look and later brought back friends for more viewing. I brought over my digital camera and 35mm Panoptic and took some pictures, just to see if I could get anything decent. I've only had a very few minutes even looking through an H-alpha filter set up and no experience with imaging through one. I was very surprised when I got home and put them on a regu-

lar monitor to see that they turned out pretty good. For me, that afternoon of solar viewing (and a little bit of imaging) was one of the big highlights of the star party. And it was a lot of fun to do a collaboration with my friend with the solar equipment and my binoviewer for some awesome viewing. Now all I need is a solar set up like that! Here are the details for those who would like to know:

Scope - 90mm TMB refractor with a 90mm Coronado H-alpha filter

Camera -- Olympus C-2100 digital attached to a 35 Panoptic with a TV adapter, zoomed all the way in (10x). Processed with Paint Shop Pro 7

<http://home.earthlink.net/~chikadee/prom1011011.jpg>  
<http://home.earthlink.net/~chikadee/prom1013013.jpg>  
<http://home.earthlink.net/~chikadee/prom1013015.jpg>

## About the Authors

### Solar Activity Report



The first observation by *Richard (Rik) Hill* (solar activity report) was the transit of Mercury in 1957. He started observing the planets in the early 1960s and has been a member of ALPO since 1975; he is a member of the board of directors of the ALPO. Rik worked on the Burrell Schmidt telescope (Kitt Peak

for 12 years and is currently with the Planetary Occultations Group and the Catalina Sky Survey (looking for potential Earth impactors) at the Lunar & Planetary Lab, Arizona University. In his spare time -- when not doing ALPO business -- Rik can be found at his C11 in the backyard or working with his fossils. He can be reached via e-mail at [rhill@lpl.arizona.edu](mailto:rhill@lpl.arizona.edu).

### ALPO Observations of Venus (1998)



*Dr. Julius L. Benton, Jr.* (Venus apparition report) is a native of Albany, Georgia, and joined the ALPO in 1967. He was appointed coordinator of the ALPO Lunar Section in 1970, heading up the Selected Areas Program, and in 1971, he assumed additional duties as coordinator of the Venus and Saturn Sections. He still holds all three posts

today, after nearly three decades of successful leadership and administration of ALPO observing programs. In 1994, Julius was elected to the board

of directors of the ALPO, and he has served as distribution editor for the *Journal of the ALPO* since 1996. In 1998 he became Associate Director of the organization. He is currently chairman of the ALPO board of directors. In addition to his professional research that has appeared in various technical journals, Julius has written extensively on the subjects of lunar and planetary astronomy for over 25 years. He can be reached at e-mail address: jlbaina@msn.com

### Hook-Like Shadow on Plato's Floor

Co-authored by Giancarlo Favero (Osservatorio Guido Ruggieri, Padova, Italy), Raffaello Lena (GLR, Geologic Lunar Researches, Italy), Fabio Lottero (GLR, Geologic Lunar Researches, Italy), and Marco Fiaschi (Osservatorio Giuseppe Colombo, Padova, Italy).



*Raffaello Lena* founded the Geologic Lunar Research (GLR) group (<http://digilander.iol.it/gibbidomine>) and has published lunar articles in the *JALPO*, *Selenology* and Italian magazines. His primary interest is the study of lunar domes and their classification. The GLR group has members from several nations

that participate in various lunar observing projects (domes, TLP, lunar flashes projects). Raffaello was recently named coordinator of the American Lunar Society's Lunar Impact Project. As such, he has done research of spurious flashes. He can be reached at e-mail address: gibbidomine@libero.it

*Giancarlo Favero's* (no photo supplied) interest in planetary astronomy was inspired about 1960 while he was in high school by Guido Ruggieri, a leading Italian planetary observer. With a home-made 7-inch reflector Giancarlo started observing Jupiter, Mars and the Moon. He received his degree in chemistry in 1969 and maintained his interest in astronomy, frequenting the Padua's Observatory and enjoying the friendship of his director, Prof. Leonida Rosino. Over the 1970s, Giancarlo directed the Jupiter Section of the Unione Astrofili Italiani (UAI) and published three papers in the *Journal of the Assn. of Lunar & Planetary Observers*. He was president of the UAI for six years, and for 30 years was professor of chemistry at the University of Padua. In 2000, he retired and subsequently joined the ALPO and returned to planetary astronomy, his youth's love.

No data supplied on Messrs. Lottero and Fiaschi

### Observations of the Remote Planets



*Richard Schmude Jr.* is coordinator of both the Remote Planets Section and the Jupiter Section of the Assn. of Lunar & Planetary Observers, and is a member of the organization's board of directors. He first became interested in astronomy when he was six years old. At that time he noticed what

seemed to be hundreds of stars in the sky -- this inspired him. He purchased his first telescope when he was 15 and showed a neighbor girl (Kathy) the Moon through this telescope. Since then, he has given over 200 astronomy presentations and telescope viewing sessions to the public. He is a member of the Association of Lunar and Planetary Observers (ALPO), the Royal Astronomical Society of Canada (RASC) and of the American Association of Variable Star Observers (AAVSO). His research interests include: measuring the brightness and color of the Moon and planets, planetary nebulae and the impact of acid rain on the environment. Richard lives in Barnesville, Georgia.

### Index to *Strolling Astronomer* Volume 42



*Michael Mattei* joined the ALPO in 1963 and has been observing the Moon and planets regularly ever since then. His professional career began at Harvard Observatory in 1967; since then, he has worked at the Wallace Astrophysical Observatory, Lincoln Labs, both of MIT, and now in the field of optics. He currently works as a senior

optical technician at Schlumberger Silicon Division. Mike continues to build telescopes and optics at home, and do CCD and video imaging of the Moon and planets with a 6-inch Schupmann telescope which he built in 1983. He is currently working on a 10-inch Schupman.

### Errata



Frank Melillo's photo was inadvertently placed next to the writeup of another author. This is Frank Melillo. Remember him. You may meet him and need to know him. We regret the error.

## ALPO Feature: Solar Activity Report Carrington Rotations 1911-1922 (1996-06-28.70 to 1997-05-22.10)

By Richard E. Hill  
ALPO Solar Section Coordinator

Solar activity was low to nonexistent with only 69 numbered active regions on the Sun during this time. The average R(I) was 9.8 and R(A) was 9.7, very similar to the last report period (Carrington Rotations 1901-1910). All but one of the rotations had a number of days with a count of zero which meant that the variations in the averages were largely determined by the highs or bulk of the days with sizable numbers. The highest sunspot count for any rotation was 20.4 in CR1916 for R(I) and 19.4 in CR1921 for R(A) with the highest daily number for R(I) being 57 on 1996-11-25 (CR1916) and for R(A) 67 on 1997-05-21 (CR1922). Figure 1 is a graph of the rotational means for this reporting period. Since the official minimum occurred in May 1996, solar activity was very low. The increase is slight but note that the highest activity, whether tabulated by rotational or daily counts, was in the last half of this reporting period. There were a number of sunspot groups that exceeded 300 millionths of the disk in area (standard unit of measure for that parameter) in contrast to the last report where none exceeded that value. One even attained an area of 1000 millionths.

As one might expect, there were few observers and observations covering this period but still more than in the last report. This is not surprising since it was just past solar minimum. However it is again worth saying that observations are as important at low activity as at high. This is how solar minimum can be determined and understood.

Terms and abbreviations used in this report are defined in the newly revised Astronomical League book "Observe and Understand the Sun". Details on obtaining this

can be found on the Solar Section webpage at:

<http://www.lpl.arizona.edu/~rhill/solar.html>

Sunspot classifications used in this report are in the McIntosh Classification System also explained in the new Astronomical League publication "A Three-Dimensional Sunspot Classification System". This is reprinted from an article in the *Strolling Astronomer* (*Journal of the Assn. of Lunar & Planetary Observers*) 33, Nos. 1-3, Jan., 1989, pp.10-13.

Times used in this report are Universal Time (U.T.). Directions are abbreviated (e.g. , N, E, SW etc.) and are celestial. Positional angular dimensions are heliographic while sunspot dimensions may be expressed in celestial arc dimensions (sec., min., etc). The term "group" refers to white-light collections of sunspots, while the term "region" refers to whole areas of activity in all wavelengths. All areas will be expressed in the standard millionths of the visible solar disk (500 millionths is about the minimum area needed for a keen-eyed observer to see a naked eye sunspot). Active Regions and sunspot groups are designated with the prefix "AR" and are designated by the Space Environment Center (SEC) of the National Oceanic and Atmospheric Administration (NOAA) in Boulder, Colorado.




Figure 1: R(I) and R(A) for rotations 1911-1922 (1996-06-28.70 to 1997-05-22.10)

**Table 1: Observers in Alphabetical Order**

Telescope					
Name	Aperture (cm)	F.L. (cm)	Type (cm)	Stop	Location
Brian Cudnik	11.3	90	rfl	n/a	Houston, TX
Michael DeTraglia	8	90	rfr	n/a	Wasilla, AK
Gordon Garcia	13	102	rfr	n/a	Hoffman Estates, IL
Richard Hill	12.5	125	s-c	n/a	Tucson, AZ
Paul Maxson	25	152	rfl	15	Phoenix, AZ
Anthony Portoni	40	163	rfl	23	Little Rock, AR
Jeffery Sandel	6	70	rfr	n/a	Cayce, SC
Fan-Lin Tao	25	375	rfr	n/a	Taipei, R.O.C.
Brad Timerson	11.4	127	rfl	n/a	Newark, NY

There were 9 observers for this reporting period.

**Table 2: Solar Cycle Rotation 23, Rotation 1911**

DATES: from 1996 06 28.70 to 1996 07 25.90					
	Mean	Max. & Date		Min. & Date	
R(I)	7.6	27	7/09	0	14 days
R(A)	7.4	28	7/09	0	14 days

This reporting period opens with activity at extremely low levels, the least since CR 1905, with only three Active Regions (ARs) designated in this rotation though the sunspot numbers themselves did not reflect this.

As this rotation began there was a decaying region on the Sun, AR7973. The beginning of its passage was covered in the last report. On 06/19 of the last report (and rotation) Maxson, at 1523UT, showed the region to consist of a single round spot with a symmetrical penumbra, class Hsx. It remained this way until it left the disk on 07/01 but was photographed by Garcia in two excellent white light images on 06/29 at 1404 and again at 1437UT. These showed a large spot surrounded by a filigree of faculae, part of it forming a ring around the spot. In an H-alpha image by Garcia the next day (1538UT) the region looked unremarkable.

Another region, AR7978, was notable in this rotation. It was first recorded by Maxson in a white light photograph on 07/07 (1632UT) just past the central meridian (CM). At that time it consisted of a spot with rudimentary penumbra, a collection of pores preceding and fol-

lowing. This observation was made at exactly the same time as the Space Environment Center produced one of their global active region maps that indicated there were no groups on the Sun!

The next day Timerson (1424UT), in a white light image, found significant development. The area was about 200 millionths of the disk and the class was Dai with a leader that was a few umbrae in penumbra with a trail of umbrae in rudimentary penumbrae leading back to the middle spot. The middle spot was a collection of six umbrae in penumbra and the follower spot was a collection of small umbrae

aligned N-S (north-south) each with its own rudimentary penumbra. Sandel, in a white light drawing on 07/09 (1830-1903.2UT) showed the leader and the trail of umbrae to be merging into one penumbra. The middle spot was just two very small collections of small umbrae each in a rudimentary penumbra aligned N-S to each other. There were small pores scattered in the middle region as well. The follower or eastern most spot, was in three pieces each with several umbrae in rudimentary penumbra. The area by this time had doubled and it was generally classed as Eki. Again on 07/10 Sandel caught this group in a whole disk drawing (1915-35UT), as it neared the limb. The area was reduced to 350 millionths and the class was Eho. The leader was now more round with two large umbrae and the middle was just a collection of tiny umbrae and pores. The follower was a single spot of three umbrae in one penumbra.

As this group was at the limb it was clearly in decay. The area was now down to 220 millionths. In a Timerson white light photo of 07/11 (1528UT) and another Sandel whole disk drawing (2105-25UT) the leader was shown to be one large umbra in a symmetrical penumbra. The middle spot was reduced to two tiny umbrae and a collection of pores. The follower was two spots with penumbrae and appeared as one spot that may have been bisected by a light bridge. On the last day of visibility for the group Sandel, again in a whole disk drawing (2105-25UT), observed that the group was reduced to just a leader that consisted of a single spot on the limb followed by two tiny umbrae. As when it came onto the disk, there were a lot of attendant faculae.

**Table 3: Solar Cycle 23, Rotation 1912**

DATES: from 1996 07 25.90 to 1996 08 22.13					
	Mean	Max. & Date	Min. & Date		
R(I)	15	23	08/02-03	8	07/26
R(A)	14	21	08/02	8	3 days

Although all levels of activity at least doubled, still the overall level of activity was very low. There were six active regions designated during this time period, exactly double that of CR1911. This was the only rotation with more than zero for minimum sunspot numbers. Only two observers contributed to this rotation.

Only one region was worth a mention during this rotation, AR7981, the remnants of old AR7978 from the previous rotation. This region was not observed by the Solar Section as it came onto the disk on 07/28 with an area already over 100 millionths, but was first seen by Sandel in a whole disk white light drawing on 08/02 (1840-55UT). At this time the leader consisted of a large umbra in a symmetrical penumbra. The follower was two small spots oriented N-S with penumbrae and a

scattering of tiny umbrae and pores. The area of this Eao group at this time had grown to 300 millionths. On 08/03 (1314UT) Garcia took a superb arc second quality photograph (Figure 2) that showed the leader to have a large umbra surrounded by a well developed, symmetrically radial penumbra. This was followed by tiny umbrae to the N and a larger umbra with rudimentary penumbra to the S. There was a N and S follower spot. The northern follower spot was a grouping of tiny umbrae with rudimentary penumbra while the southern follower had an umbra in an inverted 'V' shape open to the S with very rudimentary penumbra, where no filaments or fibrils seen at all. Sandel observed it, again in a whole disk white light drawing, about five hours later (1840-1905UT). He saw the leader to be relatively unchanged from the day before. The northern follower he saw as a spot with penumbra and the southern was a spot that had lost its penumbra with tiny umbrae and pores all around.

The next day, 08/04, was the last day of observation for this group by the Section members. Again, Garcia took a white light photo of the group (1343UT) that was a good photo but not the spectacular quality of the previous day. The leader was now more round but otherwise

pretty much the same. It was followed by just pores to the N but the spot to the S was a dark round umbra. The follower spot was just a few pores. This group was obviously in decay with its area now decreased to 250 millionths and the class now Cso.




Figure 2: AR 7981 from rotation 1912 seen in arc second resolution, taken by Gordon Garcia at 1314 UT on 3 Aug. with an Astro-Physics 130 mm aperture Starfire EDT apochromat refractor using a Baader Herschel wedge prism with an ND8x and 25a (red) filter. North is at the bottom, west to the right. Note the penumbral detail (grains and fibrils) as well as the rudimentary penumbrae that lack these structures, detached bits of penumbrae, umbral spots and pores (not as dark as the umbral spots).

**Table 4: Solar Cycle 23, Rotation 1913**

DATES: from 1996 08 22.13 to 1996 09 18.39					
	Mean	Max. & Date		Min. & Date	
R(I)	6.5	19	08/31	0	12 days
R(A)	6.2	17	08/31	0	10 days

In CR1913 activity fell back to the extremely low levels seen in the first rotation of this report. Now, as then, there were only three regions designated by SEC during this rotation.

The best observed sunspot group of this rotation was AR7986, the remains of AR7981 (which itself was old AR7978). Its coming was preceded by a facular area on the limb as seen by Portoni in a whole disk white light drawing on 08/22 (1313UT). He again observed this area on 08/23 (1333UT) when it was measured to be about 12 arc seconds in N-S extent, with an umbra and what appeared to be a symmetrical, well developed penumbra surrounded by faculae.

On 08/24 (1451UT) Garcia produced another arc second quality photograph (Figure 3) that showed the whole group to be three umbrae in surrounding penumbra. Faculae formed a ring around the spot that now had an area of 100 millionths and a class of Hsx. In an H-alpha

photo at 1659UT Garcia showed a huge filament to the N of the whole region that spanned about 120 degrees of an arc around the region. A bright plague corresponded to the faculae and extended to the E.

There were no submitted observations from 08/25-28. But on the 29th another Portoni whole disk white light drawing was noted to show that AR7986 had "darkening on N & S edges of the penumbra." The area had reduced to 50 millionths. The main spot was round with a symmetrical penumbra and several pores to the SE. A day later Portoni (1340UT) showed this situation to be unchanged but the pores were gone. The last day of observations for this region was 09/02. Again it was Portoni that made the observation with one of his drawings. The group was now near the limb and much reduced (40 millionths) and was just an umbra surrounded by a ring of faculae with more faculae following. This was the last that was seen of this region that had lasted for three rotations.

**Table 5: Solar Cycle 23, Rotation 1914**

DATES: from 1996 09 18.39 to 1996 10 15.67					
	Mean	Max. & Date		Min. & Date	
R(I)	0	0	--	0	--
R(A)	0	0	--	0	--




Figure 3: AR 7986 from rotation 1913 seen on the limb, taken by Gordon Garcia at 1451 UT on 24 Aug. with an Astro-Physics 130 mm aperture Starfire EDT apochromat refractor using a Baader Herschel wedge prism with an ND8x and 25a (red) filter. North is at the bottom, west to the right. Note the extensive faculae.

This rotation marked the low of this reporting period. There were, as the numbers show, no sunspots that lasted more than an hour or so. Only six observations total were submitted, all by DeTraglia, and all confirmed the numbers above.

This rotation marked a unique event in the near 20 year history of the Solar Section. This is the first time since CR1732 of a totally zero sunspot count. In fact, this is the lowest count that is possible. See the end of this report for further analysis.

**Table 6: Solar Cycle, Rotation 1915**

DATES: from 1996 10 15.67 to 1996 11 19.97					
	Mean	Max. & Date		Min. & Date	
R(I)	2.3	14	11/11	0	21 days
R(A)	2.6	14	10/26 & 11/11	0	19 days

In this rotation activity was elevated to the lofty heights of 'extremely low'. Still there were four Active Regions designated by SEC/NOAA; though the activity increased, the observations did the opposite. Only one observation was submitted in this rotation, by DeTraglia and it showed no spots on that day.

**Table 7: Solar Cycle, Rotation 1916**

DATES: from 1996 11 11.97 to 1996 12.09.28					
	Mean	Max. & Date		Min. & Date	
R(I)	20.4	57	11/25	0	6 days
R(A)	18.6	50	11/25-26	0	12/03 & 06

In only two rotations we jump from the lowest activity in this report to the maximum. Though this was the highest activity, still it only attained the category of 'very low'. Nine Active Regions were designated and, of these nine, there were only two observations of just one of the Active Regions, both Portoni whole disk white light drawings of AR7999.

On 11/26 Portoni, at 1411-25UT, observed this group, now with an area of 900-1000 millionths and class of Eki. This was easily the most developed group of this reporting period. On this day the leader was 4-5 umbrae in a comma-shaped penumbra with a curl to the E on the S end. In the middle of the group were three umbrae in a penumbra (possibly rudimentary), and the follower was six umbrae in one penumbra. There were some additional small umbral spots between the middle and follower spots.

In the second observation on the next day, Portoni showed that the group was smaller (with an area of 800-900 millionths) now an Ekc class. The leader was one large and three small umbrae in a single penumbra followed by the cut off tail of the comma shape or former middle spots containing 3-4 tiny umbrae. The follower was a cluster of three collections of small umbrae in rudimentary penumbrae. Clearly the group was decaying.

**Table 8: Solar Cycle 23, Rotation 1917**

DATES: from 1996 12 09.28 to 1997 01 05.61					
	Mean	Max. & Date		Min. & Date	
R(I)	13.8	22	12/13 & 18	0	6 days
R(A)	12.6	26	12/13 & 14	0	6 days

Activity in this rotation fell back from the high of the last rotation. Still categorized as 'very low' there were nevertheless six Active Regions designated but only two over 100 millionths. Observations by the Section fell to the lowest level thus far with no observations being submitted at all.

**Table 9: Solar Cycle 23, Rotation 1918**

DATES: from 1997 01 05.61 to 02 01.95					
	Mean	Max. & Date		Min. & Date	
R(I)	6.2	17	01/16	0	10 days
R(A)	6.3	16	01/16	0	7 days

In this rotation, activity fell even further to half of CR1917 and back into the 'extremely low' category. Again six Active Regions were designated, none over 50 millionths. Observers maintained the level of the last rotation as no observations were submitted.

**Table 10: Solar Cycle 23, Rotation 1919**

DATES: from 1997 02 01.95 to 1997 03 01.29					
	Mean	Max. & Date		Min. & Date	
R(I)	7.5	31	02/04	0	13 days
R(A)	7.9	27	02/03	0	9 days

Levels of activity were only slightly increased from the last rotation, but still extremely low. Four active regions were designated but all were less than 100 millionths in area and short lived. As with the last two rotations there were no observations submitted.

**Table 11: Solar Cycle 23, Rotation 1920**

DATES: from 1997 03 01.29 to 1997 03 28.60					
	Mean	Max. & Date		Min. & Date	
R(I)	7.7	25	03/15	0	12 days
R(A)	7.0	18	03/14	0	11 days

In CR1920 activity continued at extremely low levels. There were a few more regions designated, a total of seven, by SEC. The largest of these regions was AR8020 which managed to attain a maximum area of 150-200 millionths. Only one observer submitted observations, DeTraglia. Of his seven whole disk white light drawings, only three covered this sunspot group.

The first observation was on 03/10 (0030UT) when the region was already in decay with an area of only 50 millionths and a class of Cao. The leader was elongated N-S and consisted of two to three umbrae in penumbra only on the W and S sides. This was followed by a lone umbra. The next day the group had shrunk to 40 millionths but was still Cao. A DeTraglia drawing (2250UT) showed the leader as two umbrae with no penumbra followed by fragmentary penumbral bits. Two days later, 03/13 (2215UT) it was half as big and was only a collection of 8-9 umbrae with no penumbrae at all.

**Table 12: Solar Cycle 23, Rotation 1921**

DATES: from 1997 03 28.60 to 04 24.87					
	Mean	Max. & Date		Min. & Date	
R(I)	17.2	37	04/03	0	04/20-22
R(A)	19.4	43	04/03	0	04/20-21

The activity in this rotation rose to very low levels. There were nine Active Regions of which only two were observed by Section members and the observations were so sparse as to only provide a snapshot.

On 03/31 Garcia, in a region photograph (1707UT) and a whole disk white light photograph (1734UT), captured A8026 in decay and at this time only 80 millionths in area and class Dao. The leader was two spots, each with an umbra contained in rudimentary and fragmentary penumbrae. The follower spot was made up of five umbrae in rudimentary penumbra in a comma shape. A day later an H-alpha, whole disk image by Garcia showed the region to be fairly quiescent. The area was now 70 millionths but still class Dao.

AR8031 was first observed by Garcia on 04/13 in a region photograph at 1715UT of 2-3 arc seconds resolution. The leader was a line of umbrae with penumbra only to the N and S. In a middle region were pores and a few tiny umbral spots. The follower spot consisted of three to four umbrae in a symmetrical penumbra with one tongue of umbrae to the west that reached across the penumbra to the photosphere. Across the region from

the middle umbrae to the follower spot were bits of fragmentary penumbrae. The area of this Dso group was 60 millionths. In an H-alpha, whole disk photo, again by Garcia, a bright region was seen between these three spot collections but otherwise the region was unremarkable.

**Table 13: Solar Cycle 23, Rotation 1922**

DATES: from 1997 04 24.88 to 1997 5 22.10					
	Mean	Max. & Date		Min. & Date	
R(I)	13.3	52	05/21	0	4 days
R(A)	14.8	69	05/21	0	05/01

Solar activity, though still low, fell back a bit in this closing rotation of this report. Even so, the number of Active Regions rose to ten. Participation by observers was also still low and we only have one observation of AR8040, the largest of this rotation.

On 05/21 Garcia, in a region photograph (1325UT) and a whole disk white light image (1358UT) showed the leader to be an E-W elongated, teardrop shaped umbra with radial penumbra. The tip of the teardrop cut off to the E. In the middle were a few pores with a follower to the E that was a collection of about a dozen small umbrae in rudimentary penumbra aligned in a right angle that was pointed N and W. The classification at this time was Cso with 100 millionths area.

### Conclusion

This report covered a period during and just after sunspot minimum and the change over from Cycle 22 to 23. Activity was low to extremely low with those sunspot groups that did develop being of classes A-D and only one making it to E class. Unfortunately, observations decreased more than the activity itself. Even though it was minimum, there was still plenty of activity to observe, save for one rotation. More observations would have helped our understanding of the evolution of these sunspot groups. Perhaps, with better coverage, CR1914 would not have been completely spotless. Sunspot groups of modified Zurich Class (in the McIntosh system) of A and B can spring up and vanish in a matter of an hour or so. So these require vigilance and perseverance.

But readers take heart! As this is being written, we have just passed solar maximum with record numbers of observations and you can be assured that the situation in this report is only temporary.

### References

For computation of R(I) numbers, positions, classifications and designations formerly the publication Solar-Geophysical Data (Prompt Reports), Part I, was used. As of this report, it has been replaced by downloaded data from the National Geophysical Data Center at: <http://www.ngdc.noaa.gov/stp/SOLAR/getdata.html> No other references were needed since even more was there than in the published hardcopy reports!

## ALPO Feature: ALPO Observations of Venus During the 1998 Western (Morning) Apparition

By Julius L. Benton, Jr.,  
Coordinator, ALPO Venus Section

### Abstract

This report is a concise summary of photo-visual observations contributed to the ALPO Venus Section by observers in the United States, Italy, and Germany during the 1998 Western (Morning) Apparition, including instrumentation and data resources used in making those observations. Comparative studies deal with observers, instruments, and visual and photographic data. The report includes illustrations and a statistical analysis of the categories of features in the atmosphere of Venus, including cusps, cusp-caps, and cusp-bands, seen or suspected at visual wavelengths, both in integrated light and with color filters. Terminator irregularities and the apparent phase are discussed, as well as coverage based on results from continued monitoring of the dark hemisphere of Venus for the Ashen Light.

### Introduction

A total of 153 drawings and photographs of Venus was received by the ALPO Venus Section during the 1998 Western (Morning) Apparition. Geocentric phenomena in Universal Time (UT) for the Apparition are presented in *Table 1* (page 17), while *Figure 1* (page 18) illustrates the distribution of observations by month during the observing season.

**Table 1: Geocentric Phenomena in Universal Time (UT) for the 1998 Western (Morning) Apparition of Venus**

Inferior Conjunction	1998 Jan	16d 11h UT
Initial Observation	Jan	17 13
Greatest Brilliancy ( $m_v = -4.6$ )	Feb	20 02
Greatest Elongation West ( $46.5^\circ$ )	Mar	27 19
Dichotomy (predicted)	Mar	28 16
Final Observation	Oct	24 19
Superior Conjunction	Oct	30 04
<i>Apparent Diameter (observed range):</i> 62".47 (1998 Jan 17) - 9".73 (1998 Oct 24) <i>Phase</i>		
<i>Coefficient, k (observed range):</i> 0.005 (1998 Jan 17) - 0.999 (1998 Oct 24)		

ALPO observers monitored Venus reasonably well throughout the 1998 Western (Morning) Apparition, starting their observing programs just one day after Inferior Conjunction (which occurred on 1998 Jan 16) and continuing up to six days before Superior Conjunction on 1998 OCT 30. Such a consistent coverage of Venus throughout an apparition is a welcome trend that has been occurring in recent years. The "observing season," or observation period, ranged from 1998 Jan 17 to 1998 OCT 24, with over seven-eighths of the observations (88.2 percent) submitted for the period from 1998 February through August. In particular, the observations sharply peaked in 1998 February and March, the months in which Venus reached greatest brilliancy and maximum elongation from the Sun, respectively.

Seven individuals contributed visual and photographic observations of Venus during the 1998 Western (Morning) Apparition, and *Table 2* (page 18) gives their observing sites, number of observations, and instruments used.

*Figure 2* (page 19) shows the distribution of observers and contributed observations by nationality of observer for the 1998 Western (Morning) Apparition. Nearly three-quarters of the individuals taking part in ALPO Venus programs (71.4 percent) resided in the United States and accounted for slightly more than half (51.0 percent) of the total observations received. Thus, during 1998, as in recent previous apparitions, international participation in our programs continued, supporting our efforts to foster increased cooperation among lunar and planetary observers worldwide.

The types of telescopes employed to perform observations of Venus in the 1998 Western Apparition are shown in the graph in *Figure 3* (page 19). It can be seen that the great majority (97.4 percent) of observations were made with telescopes 15.2 cm (6.0 in) in aperture or greater. Classical designs (refractors and Newtonians) were used in slightly over half (52.3 percent) of the observations, while most of the remainder (47.7 percent) of Venus reports were generated using Schmidt-Cassegrains, with one observation made using a Maksutov telescope. During the 1998 Western Apparition, nearly three-fourths of the observations (74.5 percent) occurred under dark sky conditions, with the remainder occurring in twilight. Some observers followed Venus after sunrise into broad daylight in an attempt to minimize the effects of overwhelming glare associated with the planet. Also, viewing Venus higher in the sky helped reduce the effects of atmospheric dispersion and image




Figure 1. Number of Observations by Month, 1998 Western (Morning) Apparition of Venus.

observations in the form of drawings appear in this report to assist the reader in interpreting the phenomena reported in the atmosphere of Venus in the 1998 Morning Apparition. (See Figures 6-11 (pp. 21 and 22).

The visual and photographic data for the 1998 Western Apparition represented all of the traditional categories of dusky and bright markings in Venus's atmosphere (with the exception of radial dusky features), as described in the literature listed at the end of this report. *Figure 4* (page 20) shows the frequency that

distortion near the horizon, at least until late morning when solar heating caused image deterioration.

The ALPO Venus Section Coordinator extends his gratitude to the seven individuals mentioned in this report for their perseverance during early morning hours to study the planet and submit observations during the 1998 Western Apparition. Readers interested in finding out more about the Venus programs are urged to join the ALPO and become regular contributors to our observational pursuits in forthcoming apparitions.

the specific forms of markings were seen or suspected. Most observations referred to more than just one category of marking or feature, and consequently totals exceeding 100 percent can occur. Readers should recognize that there is an inherent subjectivity that exists when visual observers attempt to try to describe the very elusive atmospheric markings of Venus, and this probably affected the data in *Figure 4*. Even so, the conclusions deduced from these data appear reasonable.

The dusky markings in the atmosphere of Venus are notoriously hard to see visually, a characteristic of the

### Observations of Venusian Atmospheric Details

The standard methods and techniques for conducting visual studies of the vague and elusive "markings" in the atmosphere of Venus are outlined in detail in *The Venus Handbook* (Benton 1987). Readers who have access to earlier issues of this *Journal* may also find it helpful to consult previous apparition reports for a historical perspective on ALPO studies of Venus.

The observations used in this analysis were all made at visual wavelengths, and six examples of these

Table 2: Participants in the ALPO Venus Observing Program During the 1998 Western (Morning) Apparition

Observer and Observing Site	Number of Observations	Telescope(s) Used*
Benton, Julius L., Jr.; Wilmington Island, GA	46	15.2-cm (6.0-in) REF
Boisclair, Norman J.; South Glens Falls, NY	1	8.9-cm (3.5-in) MAK
	1	20.3-cm (8.0-in) NEW
Frassati, Mario; Crescentino, Italy	10	20.3-cm (8.0-in) SCT
Haas, Walter H.; Las Cruces, NM	2	15.2-cm (6.0-in) NEW
	8	20.3-cm (8.0-in) NEW
Niechoy, Detlev; Göttingen, Germany	1	6.0-cm (2.4-in) REF
	60	20.3-cm (8.0-in) SCT
	4	30.5-cm (12.0-in) NEW
Post, Cecil; Las Cruces, NM	4	15.2-cm (6.0-in) NEW
	12	20.3-cm (8.0-in) NEW
Venable, Roger; Augusta, GA	2	12.8-cm (5.0-in) SCT
Total Number of Observers	7	
Total Number of Observations	153	

\* MAK = Maksutov, NEW = Newtonian, REF = Refractor, SCT = Schmidt-Cassegrain.




Figure 2. Number of Observations and Observers by Nationality, 1998 Western (Morning) Apparition of Venus.

planet that is mostly independent of the experience of the observer. Using color filters and variable-density polarizers helps reveal subtle cloud phenomena on Venus at visual wavelengths, but the ALPO Venus Section strongly encourages observers to attempt regular UV (ultraviolet) photography. The morphology of features revealed at UV wavelengths is typically different from what is seen in visual regions of the spectrum, particularly with the radial dusky patterns.

Figure 4 shows that about one-third (32.7 percent) of the observations of Venus in the 1998 Western Apparition referred to a brilliant disk completely devoid of markings. When dusky features were seen or suspected, most fell in the categories of “Banded Dusky Markings” (46.9 percent), “Amorphous Dusky Markings” (45.1 percent), and “Irregular Dusky Markings” (47.8 percent). There were no sightings of “Radial Dusky Markings” during the 1998 Western (Morning) Apparition.

Terminator shading was apparent during much of the 1998 observing season, reported in 75.8 percent of the observations, as shown in Figure 4. The terminator shading usually extended from one cusp region to the other, and the shading appeared to lighten (i.e., take on a higher inten-

sity) as one progressed from the region of the terminator toward the bright limb of the planet. This gradual variance in brightness ended in the Bright Limb Band in most accounts. No photographs in the 1998 Western Apparition clearly showed any hint of terminator shading.

The mean relative intensity for all of the dusky features on Venus in the 1998 Western Apparition ranged from 8.0 to 8.5, where 0.0 represents black and 10.0 the brightest possible features. The ALPO Scale of Conspicuousness

(which runs sequentially from 0.0 for “definitely not seen” up to 10.0 for “certainly seen”) was also used regularly during this apparition. On this scale, the dusky markings in Figure 4 had a mean conspicuousness of about 3.0 during the apparition, which suggests that these features fell within the range between very indistinct impressions and fairly good indications of their actual presence on Venus.

Figure 4 also shows that “Bright Spots or Regions,” exclusive of the cusp areas, were not seen or suspected in any of the submitted observations. When seen, it is customary in drawings made at visual wavelengths for




Figure 3. Number of Observations by Type of Telescope, 1998 Western (Morning) Apparition of Venus.




Figure 4. Relative Frequency of Specific Forms of Atmospheric Markings, 1998 Western (Morning) Apparition of Venus.

observers to call attention to such bright areas by sketching dotted lines around such features.

Observers regularly used color-filter techniques during the 1998 Western (Morning) Apparition, and when results were compared with studies in Integrated Light (no filter), it was clear that color filters and variable-density polarizers helped improve the visibility of elusive atmospheric phenomena on Venus.

### The Bright Limb Band

Figure 4 reveals that 58.8 percent of the submitted observations in the 1998 Western Apparition reported a “Bright Limb Band” on Venus’s sunlit hemisphere.

When the Bright Limb Band was reported, it appeared as a continuous, brilliant arc extending from cusp to cusp for only 28.6 percent of the observations, and was interrupted or only partially visible along the limb of Venus in 71.4 percent of the positive reports. This impression by observers in 1998 is in sharp contrast with other recent apparition reports. The mean numerical intensity of the Bright Limb Band was 9.7, becoming more obvious when color filters or variable-density polarizers were employed. Despite the

dazzling brilliance of this feature for visual observers, it was not apparent in any photographs of Venus submitted in 1998.

### Terminator Irregularities

The terminator is the geometric curve that divides the sunlit and dark hemispheres of Venus. Observers described an irregular or asymmetric terminator in a little less than one-third (30.1 percent) of the observations in the 1998 Western Apparition. Amorphous, banded, and irregular dusky atmo-

spheric markings appeared to blend with the shading along the terminator, possibly contributing to reported deformities. Filter techniques enhanced the visibility of terminator irregularities, and dusky atmospheric features closely associated with them, during this apparition. Because of irradiation, bright features adjacent to the terminator may occasionally look like bulges, and dark features may look like dusky hollows.

### Cusps, Cusp-Caps and Cusp-Bands

In most cases, when the *phase coefficient*,  $k$ , the fraction of the disc that is illuminated, lies between 0.1 and 0.8,




Figure 5. Relative Frequency of Observations of Specific Cusp Features, 1998 Western (Morning) Apparition of Venus.

NOTE: The six drawings that follow are all oriented with celestial south at the top and celestial west at the left, which is the normal inverted view when observing objects near the meridian with an astronomical telescope in the Earth's Northern Hemisphere. (Figures 6 and 7 were originally reversed and have been rectified.) Seeing was reported, or has been converted to, the standard ALPO scale, ranging from 0.0 for the worst possible condition to 10.0 for perfect seeing. Transparency is on the ALPO Scale, ranging from 0 for worst to 5 for perfect.




Figure 6. 1998 Feb 02, 11:43 UT. Detlev Niechoy, Göttingen, Germany. 20.3-cm (8.0-in) Schmidt-Cassegrain, 51X, Integrated Light. Seeing 2.5, Transparency 2.5. Ashen Light definitely visible. Phase ( $k$ ) = 0.098, Diameter = 53".5.




Figure 7. 1998 Mar 26, 05:46 UT. Detlev Niechoy, Göttingen, Germany. 20.3-cm (8.0-in) Schmidt-Cassegrain, 112X, Integrated Light. Seeing 2.0, Transparency 2.0. Phase ( $k$ ) = 0.486, Diameter = 25".0.

features on Venus having the most contrast and prominence are repeatedly sighted at or near the planet's cusps. These cusp-caps are sometimes bordered by what are described as dark, usually diffuse, cusp-bands. Figure 5 (page 20) shows the visibility statistics for Venusian cusp features in the 1998 Western Apparition.

Figure 5 shows that the northern and southern cusp-caps of Venus, when observed, were equal in size and brightness most of the time. There were a very few instances when either the northern or southern cusp-cap was the larger, the brighter, or was both, and in slightly more than one-third of the observations submitted (36.6 percent), neither cusp-cap was visible. The mean relative intensity of the cusp-caps was about 9.9 during the 1998 Western Apparition. Dusky cusp-bands bordering the bright cusp-caps were not reported in 41.1 percent of the observations when cusp-caps were visible, and the cusp-bands displayed a mean relative intensity of about 6.8 (see Figure 5).

## Cusp Extensions

As shown in Figure 5, there were no cusp extensions reported beyond the 180° expected from simple geometry for 92.8 percent of the observations (both in integrated light and with color filters). Early in the apparition, as Venus progressed through its crescentic phases following inferior conjunction in 1998, several observers recorded cusp extensions that ranged from 2° to 15°. Just after inferior conjunction, a few observers thought that the cusps joined along the planet's unilluminated limb, forming a beautiful halo encircling the dark hemisphere of Venus. Reported cusp extensions were shown on drawings, with their appearance enhanced by color filters and polarizers, but none were photographed successfully. Experience has shown that cusp extensions are very difficult to document on film due to the fact that the sunlit regions of Venus are so much brighter than the faint extensions. We encourage observers to try recording cusp extensions with CCD or video cameras in future apparitions.

## Estimates of Dichotomy

A discrepancy between the predicted and the observed dates of dichotomy (half-phase), known as the "Schroeter Effect" on Venus, was not reported by observers during the 1998 Western (Morning) Apparition as no observers submitted phase estimates during the apparition. The predicted half-phase occurs when  $k = 0.500$ , and the *phase angle*,  $i$ , between the Sun and the Earth as seen from Venus equals 90°.

## Dark-Hemisphere Phenomena and Ashen-Light Observations

The Ashen Light, first reported by G. Riccioli in 1643, refers to an extremely elusive, faint illumination of

Venus's dark hemisphere. Although it does not have the same origin, the Ashen Light resembles Earthshine on the dark portion of the Moon. Most observers agree that Venus must be viewed against a completely dark sky for the Ashen Light to be seen, but such circumstances occur only when the planet is very low in the sky where adverse terrestrial atmospheric conditions contribute to




Figure 8. 1998 Mar 29, 08:10 UT. Mario Frassati, Crescentino, Italy. 20.3-cm (8.0-in) Schmidt-Cassegrain, 222X, W8 (light yellow) Filter. Seeing 2.0 (interpolated). Terminator irregularities; cloud extending over terminator? Phase (k) = 0.502, Diameter = 24".1.




Figure 9. 1998 June 16, 10:00 UT. Mario Frassati, Crescentino, Italy. 20.3-cm (8.0-in) Schmidt-Cassegrain, 250X, W80A (light blue) Filter. Seeing 6.5 (interpolated). Phase (k) = 0.802, Diameter = 13".1.




Figure 10. 1998 July 10, 08:25 UT. Mario Frassati, Crescentino, Italy. 20.3-cm (8.0-in) Schmidt-Cassegrain, 250X, W80A (light blue) Filter. Seeing 7.0 (interpolated). Phase (k) = 0.865, Diameter = 11".8.




Figure 11. 1998 AUG 25, 08:32 UT. Mario Frassati, Crescentino, Italy. 20.3-cm (8.0-in) Schmidt-Cassegrain, 250X, W80A (light blue) Filter. Seeing 6.0 (interpolated.) Phase (k) = 0.954, Diameter = 10".3.

poor seeing. Also, substantial glare in contrast with the surrounding dark sky influences such observations. Even so, the ALPO Venus Section continues to hear from observers who say they have seen the Ashen Light when Venus was against a twilight sky.

During the 1998 Western Apparition, there were practically no instances (2.6 percent of the observations) when the Ashen Light was seen or suspected in Inte-

grated Light, or with color filters or variable-density polarizers. On 1998 Jan 31, 11:34 to 11:40 UT, veteran Venus observer Detlev Niechoy of Göttingen, Germany reported that the Ashen Light was definitely visible in blue light and in Integrated Light (no filter) using his 20.3-cm (8.0-in) Schmidt-Cassegrain at 112X, and he described the Ashen Light as also “definite” on 1998 Feb 02, 11:43 to 12:02 UT, in Integrated Light and with a blue filter but not with W25 (red), W15 (yellow), and W47 (dark blue) filters. No other observers reported any suspicions of the phenomenon, and unfortunately, there were no confirmations of the two Ashen Light sightings by Niechoy.

Indeed, there were a few instances during the 1998 Western (Morning) Apparition (7.2 percent) when observers described the dark hemisphere of Venus as looking *darker* than the background sky, this phenomenon is almost surely an effect of contrast.

## Conclusions

The results of our analysis of visual and photographic observations contributed to the ALPO Venus Section during the 1998 Western (Morning) Apparition of Venus indicated limited activity in the atmosphere of Venus. It was mentioned earlier in this report that it is very troublesome to differentiate between what constitutes real atmospheric phenomena and what is merely illusory on Venus at visual wavelengths. The level of confidence in our results will improve as the number of observers and incidence of simultaneous observations increase. The Venus Section is making a special effort to organize and implement simultaneous observation schedules so that observers near each other can set aside mutual times to jointly follow Venus using similar methods and equipment. The simultaneous observing schedule is expected to appear on the ALPO Website in the very near future at [www.lpl.arizona.edu/alpo](http://www.lpl.arizona.edu/alpo). In addition to routine observations, the Venus Section desperately needs more ultraviolet photographs of Venus, as well as CCD images of the planet at different wavelengths. We are attempting to standardize and improve observational techniques and methodology so that comparison of our results with those of previous morning observing seasons, as well as with evening apparitions of Venus, is more reliable.

ALPO studies of the Ashen Light, which peaked during the Pioneer Venus Orbiter Project, are continuing every apparition. Constant monitoring of the planet for the presence of this phenomenon by a large number of observers (ideally participating in a simultaneous observing program) remains important as a means of improving our chances of capturing confirmed dark-hemisphere events.

Active international cooperation by individuals making regular systematic, simultaneous observations of Venus remains our main objective, and the ALPO Venus Sec-

tion invites interested readers to join us in our projects and challenges ahead.

## References

- Benton, Julius L., Jr. (1973). *An Introduction to Observing Venus*. Savannah, GA: Review Publishing Company.
- \_\_\_\_\_. (1987). *Visual Observations of Venus: Theory and Methods (The ALPO Venus Handbook)*. Savannah, GA: Review Publishing Company.
- \_\_\_\_\_. (1998). “The ALPO Observations of Venus: The 1994 Eastern (Evening) Apparition.” *JALPO*, 40, No. 2 (Apr.): 54-61.
- \_\_\_\_\_. (1999). “ALPO Observations of Venus During the 1995-96 Eastern (Evening) Apparition.” *JALPO*, 41, No. 2 (Apr.): 57-65.
- \_\_\_\_\_. (2000). “ALPO Observations of Venus During the 1997-98 Eastern (Evening) Apparition.” *JALPO*, 42, No. 4 (Oct.): 149-157.
- Hunten, D.M., *et al.*, eds. (1983). *Venus*. Tucson: University of Arizona Press.
- United States Naval Observatory (1997). *The Astronomical Almanac*. Washington: U.S. Government Printing Office. (Annual Publication; the 1998 edition, which was published in 1997, was used for this report.)

### **(From the ALPO Venus Section Mission Statement)**

“Visual observations of the planet Venus should be carried out in a cooperative systematic research program to maximize opportunities for useful scientific contributions. In the A.LPO Venus Section, our objective is to observe Venus on every possible clear night throughout an apparition of the planet, which runs from conjunction to subsequent conjunction with the Sun. Venus is an inferior planet, meaning that it has a smaller interior orbit to that of the Earth, and it exhibits phases just like the Moon. Because Venus is comparatively near the Sun, it is characteristically very bright, and the high albedo produces an excessive amount of glare. The rather faint and elusive markings on the disk of Venus, normally of very low contrast, become difficult to see as a result. Considerable controversy exists over the true nature of these dusky amorphous or somewhat streaky atmospheric features. It is not at all unusual for two observers, working on the same date with comparable instrumentation, to see strikingly dissimilar atmospheric phenomena on the planet.”

**ALPO Feature:**

**The Nature of the Hook-Like Shadow on Plato's Floor Observed by Wilkins and Moore in 1952**

**Part II. Simulations with a Computer and a Plasticine® Model**

**By: Giancarlo Favero<sup>a</sup>, Raffaello Lena<sup>b</sup>,  
Fabio Lottero<sup>b</sup> and Marco Fiaschi<sup>c</sup>**

<sup>a</sup> Osservatorio Guido Ruggieri, Padova, Italy

<sup>b</sup> GLR, Geologic Lunar Researches, Italy

<sup>c</sup> Osservatorio Giuseppe Colombo, Padova, Italy

## Introduction

A paper recently appeared in this Journal [1] which proposed that the hook-like shadow recorded on the floor of the lunar crater Plato by Wilkins and Moore on 1952 April 3, [2, 3], is projected by a complex and elongated hill lying on Plato's floor, at the foot of its south wall. The hill is easily visible on Lunar Orbiter IV images [1] and has coordinates  $\eta = 0.7693$ ,  $\xi = -0.0875$  (latitude  $50.29^\circ$ , longitude  $-7.87^\circ$ ). Some Italian authors [4] and an ALPO editor [5] maintained another opinion expressed by O'Connell in 1998 [6], that the hook shadow is shed by the gamma peak (peak #3 in paper [1]), the tallest of the Plato's east rim peaks.

We carried out a computer simulation of the 1952 drawings and built a Plasticine® model of Plato in order to demonstrate that:

- The hook-like shadow recorded on the Wilkins and Moore drawings is compatible with the shadow of the complex and elongated hill, as suggested in paper [1].
- The gamma peak shadow cannot appear where the hook-like shadow is reported in the drawings, regardless of the illumination conditions.
- The gamma peak shadow never displays significant curvature so it cannot be the hook shadow.

## Experimental

### A. Computer Simulation of the 1952 Shadows

The original two Plato drawings made on 1952 April 3, 21<sup>h</sup> 30<sup>m</sup> UT [2, 3] are reproduced here as *Figures 1* and *2*. Each one was scanned and digitally converted to obtain the contour of: the crater rim, the hook-like shadow, the shadow cast on Plato's floor by its east wall, the shadows of a few features lying on Plato's east wall and the shadows of a few features lying outside




Figure 1: Plato drawn by H. P. Wilkins on 1952 April 3, at 21h 30m UT, observing the Moon with the Meudon *grande lunette* of 83 cm (altitude of the Sun over Plato,  $H = 5.32^\circ$ ; azimuth of the Sun,  $A = 94.95^\circ$ ). The hook-like shadow near the south-east wall is evident. South is up, or nearly so, in all the figures.




Figure 2: Plato drawn at the same time as Figure 1 by P. Moore, observing with the same instrument. Features on Plato's floor are similar, in particular the "hook", while those outside the crater are crudely different, revealing that the authors were concerned with few features on the floor.

Plato's wall. We selected for our simulation only those features of the last two types which were identifiable in both drawings.




Figure 3: The best fit between some shadows of the Wilkins's drawing (reticulated) and the corresponding features on the Lunar Orbiter IV image IV-127-H3

The digital image of each drawing was measured on the computer screen to obtain, in units of pixels, the length of Plato's major and minor axes. Along the same two directions, identified by the same geological features, we measured the corresponding diameters of Plato on the image IV-127-H3 taken by Lunar Orbiter IV 1967 May 20, 6<sup>h</sup> 26<sup>m</sup> UT (Sun's height over Plato,  $H = 20.79^\circ$ ; Sun's azimuth,  $A = 117.47^\circ$ ). The digital image




Figure 4: The best fit between some shadows of the Moore's drawing (reticulated) and the corresponding features on the Lunar Orbiter IV image IV-127-H3

of each drawing was then stretched and rotated to give the best superimposition over the Lunar Orbiter image. It is worth noting that the hook-like shadow was not considered in these operations. The best fit obtained with this procedure is shown in *Figure 3* for the Wilkins drawing and in *Figure 4* for the Moore drawing.

### B. The Plasticine® Model

A 6 inch diameter (150 mm) 3-D model of Plato's east wall and floor (*Figure 5*), at the scale of about 1:670.000, was built by layering Plasticine® over a copy of the Lunar Orbiter photo described above.




Figure 5: The Plasticine® model built for this work. The lighting is similar to O'Connell's image [4] and to *Figure 3* in [1]:  $H = 3^\circ$ ;  $A = 95^\circ$ .

The height in kilometres of different portions of Plato's east wall was calculated with the ALPO Lunar Observer's Tool Kit software from the relevant shadow lengths measured in units of pixels on *Figure 3* of [1] and is reported in Table 1.

The height of each measured portion of Plato's east wall, resampled with the proper scale ratio, was built up with Plasticine®. The height was determined with a centesimal calliper (0.05 mm precision, corresponding to 33 m, or 106 ft, on the Moon). Each calibrated portion was then connected to the nearby ones so as to obtain shadows like those recorded on the CCD images taken under similar lighting conditions. The slope of Plato's inner wall was obtained by connecting the wall rim with the scarp base recorded on the Lunar Orbiter photo. The Moon's curvature was neglected.

**Table 1: Coordinates, shadow lengths and peak heights of peaks 1 through 9 from Figure 3b of article [1]. For some of the valleys (a-h) eta values are also reported, but not their xi values, because the position of the highest feature on the bottom of a valley can differ considerably at different sunlight inclinations.**

Peak (1,2 ...) valley (a,b ...)	eta	xi	Shadow (pixels ±1)	Height (km ± 0.03)
a	0.7683		40	1.29
1	0.7700	-0.0822	43	1.39
2	0.7715	-0.0800	47	1.52
b	0.7723		40	1.29
3	0.7724	-0.0759	65	2.10
c	0.7750		37	1.20
4	0.7800	-0.0727	46	1.49
d	0.7814		40	1.29
5	0.7830	-0.0719	49	1.58
6	0.7859	-0.0716	53	1.71
e	0.7886		30	0.97
7	0.7905	-0.0729	39	1.26
f	0.7921		32	1.03
8	0.7941	-0.0748	40	1.29
g	0.7959		30	0.97
9	0.7968	-0.0769	35	1.11

**Table 2: Coordinates, shadow lengths and heights of the complex and elongated hill from measurements from figure 7 in [1].**

Peak	eta	xi	Shadow (pixels ± 1)	Height (km ± 0.03)
ch	0.7693	-0.0875	3	0.50

The white-painted model was placed in a dark room, illuminated by a beam of sunlight, and photographed with a camcorder. Different sequences were recorded modifying the Sun's height and azimuth. A few frames of suitable sequences were digitally converted and they are shown as *Figures 5-7*.

We included some key geological features on this model (*Figure 6*). First, we put the complex and elongated hill proposed in [1] as the cause of the hook-like shadow, adopting a height of 500 m (1550 ft) measured by its shadow in *Figure 7* of [1] and calculated with the ALPO *Lunar Observer's Tool Kit* software (Table 2). A few smaller hills lying on Plato's floor at the foot of its east wall were also included, to improve the shadows' contour for the Sun's height around 10°. The four craterlets reported on the 1952 drawings were also added to the model, with exaggerated wall heights. Also, we added to the model (*Figure 6*) the meridian passing through the gamma peak and a protractor arc, with its center on

the gamma peak, to indicate the Sun's azimuth, referred to the same peak, from 90° to 120° in 5° increments.

## Results and Discussion

### A. Computer elaboration of the 1952 shadows

The digital copies of the 1952 drawings were computer resampled to simulate Plato's perspective as shown in the IV-127-H3 Lunar Orbiter IV photo. Each resampled drawing was then superimposed on that photo to give the best fit between a few selected features appearing on both pertaining to Plato's wall rim and the neighboring terrain. The hook-like shadow was deliberately neglected in this procedure, so its final position was only the result of the best fit among the other selected features. The results are reported in *Figures 3* and *4* which show the shadows from Wilkins and Moore, respectively, superimposed on the Lunar Orbiter photo IV-127-H3. In both figures the hook-like shadow closely encircles the north-west base of the complex and elongated hill, as one would expect if the hill is the source of the hook, as suggested in paper [1].

### B. The Plasticine® Model

The fidelity of our Plato Plasticine® model (*Figure 5*) is demonstrated by its ability to simulate fairly well the shadows appearing in the O'Connell CCD image [6] and in *Figure 3* of paper [1]. An interesting result provided by

the model is that the complex and elongated hill, proposed in paper [1] as the source of the hook-like shadow, can actually cast a shadow of that geometry in the correct position, as *Figure 6* clearly shows (compare *Figure 6* with *Figures 1* and *2*).



**Figure 6: Plato's model illuminated like the CCD images of Figures 6 and 7 in [1]: H = 11°; A = 105°. The meridian passing through the gamma peak, the 90°-120° protractor and the hook-like shadow are indicated.**



Figure 7: The gamma peak's shadow would combine with the complex and elongated hill's shadow only for a Sun's azimuth of  $80^\circ$  and  $10^\circ$  Sun's height. Only thus the model would be similar to the 1952 drawings, but this azimuth is impossible.




Figure 8: Image IV-116-H1 of Lunar Orbiter IV taken 1967 May 19, 7<sup>h</sup> 00<sup>m</sup> UT ( $H = 15.71^\circ$ ;  $A = 109.02^\circ$ ). The gamma peak shadow is well visible, northward of the curved shadow of the complex and elongated hill (ch), which appears like the hook shadow of the 1952 drawings.

A further result concerns the gamma peak shadow (peak and shadow #3 in paper [1]). The gamma peak's shadow is the longest shadow, measured from the east wall rim to the shadow's tip, on Plato's floor under any lighting condition. It should coincide with the 1952 hook as proposed by O'Connell only for solar azimuth values lower than  $80^\circ$ . This azimuth of  $80^\circ$  is the azimuth simulated in Figure 7, where the gamma peak shadow is cast near the elongated hill in which case it appears similar to the 1952 hook.

But an azimuth of  $80^\circ$  is clearly impossible, so the gamma peak shadow can NEVER be in the place of the 1952 hook. While the Sun rises over the gamma peak, its shadow on Plato's floor shortens and shifts northward, because of the rising of Sun and because of its increasing azimuth. This is confirmed by the image IV-116-H1 of Lunar Orbiter IV taken 1967 May 19, 7h 00m UT ( $H = 15.71^\circ$ ;  $A = 109.02^\circ$ ) and reported here as Figure 8.

Figure 8 shows the gamma peak shadow in a position more northern than the 1952 hook. Moreover, Figure 8 shows the top of the complex and elongated hill near the Plato's southern wall: the top of the hill is illuminated by the Sun. The north-western base of the hill is encircled by shadows recorded by Lunar Orbiter. These shadows cannot be seen with the same resolution by an earth-based observer. Integrated by an earth-based observer, these shadows combine to give the hook.

## Why is the gamma peak shadow not reported on the 1952 drawings?

If what precedes is correct, we must search for an answer to this question.

We think that the two Plato drawings made at the telescope on 1952 April 3 by the two English observers were originally only rough and incomplete sketches of a few features upon which their attention was directed. This hypothesis is supported by their words [2, page 236]: "The shaded drawing [of Plato reported on page 234] .... may be relied upon as depicting, with accuracy, the relative size and intensities of the cratercones, spots and streaks (?) visible under that particular angle of illumination." The question mark concerns the absence of any streak from the Wilkins drawing (Figure 1), while three streaks are well evident in the Moore drawing (Figure 2). We can thus maintain that the observers recorded at the telescope only these few features, including the hook-like shadow, which appears similarly in the two drawings. Completing from memory their drawing in a later time, the two observers forgot to report the gamma peak shadow (probably confounding it with the hook) and introduced a considerable amount of rough differences in the Plato's neighborhoods,

which can be seen by comparing features between *Figure 1* and *2*.

This hypothesis is sustained by the following reconstruction of the 1952 April 3 minimal observing schedule based on “The Moon” [2] even though a precise time scale is lacking:

- Wilkins “saw a distinct craterlet on the summit of the central mountain [of Alpetragius], surrounded by four almost overhanging peaks. This was at once confirmed by Moore... (page 138).
- Wilkins “critically examined Conon...”. It follows a detailed description of the observed features (page 94).
- Moore ... noted ten objects on the floor [of Archimedes], only four of which appeared as distinct craterlets.” It follows a detailed description of the observed features (page 92).
- Wilkins and Moore “discovered ... a white, very shallow crater within which is a most minute central pit, the whole strongly resembling a “wash-bowl” (page 228).
- At 21<sup>h</sup> 30<sup>m</sup> UT the observers sketch Plato at the same time (page 234).
- Between 22<sup>h</sup> 00<sup>m</sup> and 22<sup>h</sup> 15<sup>m</sup> UT Wilkins draws Guerické (page 127).
- At 22<sup>h</sup> 15<sup>m</sup> UT Moore “first distinctly sees a craterlet where A and Birt join...” (page 140, the year “1951” reported on the figure caption is clearly a misprint, as confirmed by the accompanying text which states that the observation was done on “3 April 1952”).

The two observers studied Plato for not more than 30 minutes, 15 minutes each one, to cover all of those observations. It appears impossible that an observer, even an expert, could complete at the telescope a drawing as detailed as the ones in *Figure 1* or *2* in this little time.

In conclusion, we suggest that the 1952 drawings contain a number of common features which attracted the attention of both observers. These certainly include the hooked shadow, whose shape and position seem accurate under the limits of a visual observation. All other features, particularly those outside Plato but also some shadows on Plato’s floor (as we will show later), have only a vague counterpart in reality.

## Comparing the 1952 Drawings with a Similar CCD Image

*Figure 9* allows a comparison between the 1952 drawings and a CCD image (*Figure 4* in [1]) taken under nearly the same illumination conditions:  $H = 5.32^\circ$  in




Figure 9: Comparison between the 1952 drawings and a CCD image taken under similar light conditions ( $H = 5.15^\circ$ ;  $A = 98.50^\circ$ ). The CCD image shows the westward extension of the shadow south of the gamma peak’s shadow, while the shadow’s portion south of the hook is restricted toward east.

the drawings and  $5.15^\circ$  in the CCD frame;  $A = 94.95^\circ$  in the drawings and  $98.59^\circ$  in the CCD frame. The reported values are referred to the top of the gamma peak.

The shadow cast by Plato’s east wall onto its floor appears much more extended in the drawings than in the CCD image, particularly in its northern half. This fact in itself suggests a time for the drawings earlier than 21<sup>h</sup> 30<sup>m</sup> UT. However this cannot be true, because the mean height of the east wall mountains diminishes northward (Table 1). This is confirmed by all the images reported in [1], also those added by the ALPO editor. It can be concluded that the shape and extension of the shadow of the Plato east wall which appear in the 1952 drawings were not recorded at the telescope. Therefore this shadow cannot be relied upon.

As far as the hook-like shadow is concerned, *Figure 9* reveals that it appears more eastward in the drawings, particularly in the Moore one, and more southward than the gamma peak shadow in the CCD image. We are sure that the hook was recorded accurately by the two observers in their 1952 drawings, so that three conclusions may be drawn:

- The time of these drawings may be later than 21<sup>h</sup> 30<sup>m</sup> UT (according to Moore, as reported in [1]).
- The hook-like shadow cannot be identified with the gamma peak shadow, because of the different position of these two features.
- The gamma peak shadow doesn’t show any “hook” curvature in the CCD image, which has a very high quality (pixel scale about 0.8 km).

Now we wish to compare, in *Figure 9*, the portion of the shadow between the hook (or the gamma peak shadow) and Plato’s south wall. The CCD image shows that this

portion of the shadow is much extended westward, so as to largely extend beyond the longitude of the shadow tip of the gamma peak. On the opposite case, the drawings show that the shadow between the hook and the south wall is strongly receded eastward. This should confirm that the drawings were made considerably later than the time recorded, but the situation isn't so simple.

If Plato's floor was perfectly flat in the region under examination, the shadow width should have receded regularly with the Sun's rising, nearly maintaining its original western profile. In other words, the portion of shadow between the south wall and the hook (or the gamma peak shadow) should have maintained a larger extension westward of the hook-like shadow (or the gamma peak shadow). We are convinced that the strong eastward shift of the shadow south of the hook in the 1952 drawings is due to the emersion of the complex and elongated hill (height about 500 m, 1550 ft) from the east wall shadow, as shown in *Figure 8*. By now the hill casts his shadow westward originating the hook-like shadow, while the gamma peak shadow continues to be visible, more northward than the hook and progressively confused with the inner portions of Plato's east wall shadows. This happens when the Sun's height over Plato's gamma peak exceeds about  $8^\circ$ , i.e., just after 24<sup>h</sup> UT on 1952 April 3, when the Moon was about  $20^\circ$  high over the Meudon horizon.

## Conclusions

The Plato drawings made by Wilkins and Moore on 1952 April 3 show crude differences between them and a similar CCD image as far as the crater neighborhoods are concerned. To explain these differences we suggest that the English observers each executed at the telescope only a partial drawing intended only to record a few features on Plato's floor, including the hook-like shadow which appears similarly recorded by both. On this basis we try to explain also the absence from the drawings of the gamma peak shadow and the anomalous shape of the east wall shadow, which was recorded by a CCD in similar lighting conditions.

Computer simulations of the features similarly recorded on the 1952 drawings, including the hook-like shadow, allowed their comparison with the IV-127-H3 Lunar Orbiter IV image. *Figures 3* and *4* sustain the hypothesis that the hook shadow is cast by the complex and elongated hill situated on the crater floor, near its south wall, as suggested in [1]. Accordingly, we can also explain the rapid receding of the shadow comprised between the hook and Plato's south wall.

Our Plasticine<sup>®</sup> Plato model, illuminated under different Sun conditions, sustains this hypothesis and discards the involvement of the gamma peak shadow in the formation of the hook. In fact, the gamma peak shadow remains visible, but sensibly northward of the hook, in

any lighting condition from  $H = 3^\circ$  and  $A = 94^\circ$  to  $H = 11^\circ$  and  $A = 105^\circ$ , simulated by our model, and certainly up to  $H = 13.93^\circ$  and  $A = 107.27^\circ$  of the Lunar Orbiter IV image IV-116-H1.

## References

- [1] Favero, G., Lena, R., Mengoli, G., Cipolat, A. & Gualdoni, P., "The Nature of the Hook-like Shadow Observed by Wilkins and Moore on Plato's Floor in 1952", *Journal of the Assn. of Lunar & Planetary Observers*, Vol. 42, 2000, number 3, pp. 126-131.
- [2] Wilkins, H. P. & Moore, P., *The Moon*, 2nd Ed., Faber and Faber Ltd, London, 1961, p. 236 (Wilkins drawing).
- [3] Wilkins, H. P., *Our Moon*, Frederick Muller Ltd, London, 1958, p. 70 (Moore drawing).
- [4] Braga, R. & Ferri, F., "Plato Hook: l'Uncino di Plato", [http://www.uai.it/sez\\_lun/hook.htm](http://www.uai.it/sez_lun/hook.htm)
- [5] "Three Additional Views of the 'Hook' Shadow on Plato's Floor", *Journal of the Assn. of Lunar & Planetary Observers*, Vol. 42, 2000, number 3, p. 132.
- [6] O'Connell, B., "Plato Hook. Clementine and CCD images shed light on the shadowy mystery of a 45 year old drawing", Proceedings of the 49<sup>th</sup> Convention of the ALPO, Atlanta, Georgia, July 9-11, 1998, *ALPO Monograph No 8*.

## Note on Figures 1 and 2

Wilkins's and Moore's drawings (Figures 1 and 2) can be reproduced without permission from the publishers because:

1. The Faber and Faber Editor no longer holds the "The Moon" rights, as told to us by Sally Robson ([sally.robson@faber.co.uk](mailto:sally.robson@faber.co.uk)) who suggested we ask A P Watt Ltd.
2. A. P. Watt does not hold the rights of the book, as told to us by Melanie Rigg, [apw@apwatt.co.uk](mailto:apw@apwatt.co.uk)
3. The editor of the book *Our Moon*, Frederick Muller Ltd, London, is no more active following 1985.

## ALPO Feature: Observations of the Remote Planets in 2000

**Richard W. Schmude, Jr.**  
Coordinator,  
ALPO Remote Planets Section

### Abstract

Over a dozen people studied the remote planets of our solar system, Uranus, Neptune and Pluto in the year 2000 and this report summarizes those observations. The selected normalized magnitudes of Uranus are:  $B(1,0) = -6.56$ ,  $V(1,0) = -7.14 \pm 0.01$ ,  $R(1,0) = -6.82 \pm 0.01$  and  $I(1,0) = -5.62 \pm 0.05$  while the corresponding value for Neptune is:  $V(1,0) = -6.98 \pm 0.02$ . During 1991-2000, Uranus became dimmer in the V filter at an average rate of 0.0020 magnitudes per year while Neptune became 0.0112 magnitudes per year brighter in the same filter. Unfiltered CCD images show Titania to be 0.24 magnitudes brighter than Oberon on August 6, 2000. Unfiltered CCD images made in 1998 and 2000 are consistent with the Pluto system being brightest at a sub-Earth longitude of  $130^\circ$  and dimmest at  $280^\circ$ .

### Introduction

Two recent developments related to the remote planets were the detection of "clouds" on Uranus from ground-based telescopes (Sromovsky et al. 2000, 307) and the construction of a two-color map of Pluto's sub-Charon hemisphere (Young, Binzel and Crane, 2001, 552). Sromovsky and co-workers used the NASA Infrared Telescope Facility to image Uranus at several wavelengths including 1.7 microns in mid-1998 and mid-1999. They

detected faint clouds in 1998 and a bright cloud in 1999. They point out the need for further studies to determine if Uranus is about to undergo a period of increased cloud activity. Young, Binzel and Crane (2001, 552) analyzed Charon occultation data to construct a color map of Pluto. A dark feature just south of Pluto's equator was resolved and it is concluded that this feature is comprised of several distinct color units.

In addition to the two breakthroughs mentioned in the preceding paragraph, professional astronomers continue to monitor the clouds on Neptune in infrared wavelengths (Roddier et al. 1998, 168), (Talcott, 2001, 30). Wasserman, Olkin and Franz (2000, 1082) used Hubble Space Telescope images to measure a Charon/Pluto mass ratio of  $0.122 \pm 0.005$ . Finally, Frank Melillo of the Association of Lunar & Planetary Observers (ALPO) recorded methane-band images of Uranus on five different nights in 2000.

Table 1 lists the characteristics of the 2000 apparitions of Uranus, Neptune and Pluto. Sixteen people submitted observations of the remote planets between mid 2000 and early 2001; their names, locations and types of observations are listed in Table 2.

### Photoelectric Photometry

Five people submitted photoelectric magnitude measurements of the remote planets in 2000. In all cases, an SSP-3 solid-state photometer was used along with filters that were transformed to the Johnson B, V, R and I system; more information on the instrument and filters can be found elsewhere (Optec, 1997), (Schmude, 1992, 20). Katie used a 0.09 meter Maksutov telescope in

making her measurements. Both Frank and Brian used a 0.20 meter Schmidt-Cassegrain telescope in making their measurements, while John made all of his measurements with a 0.13 meter Schmidt-Cassegrain telescope. The author used a 0.51 meter Newtonian in May and June but used a 0.09 meter Maksutov telescope during September. Table 3 lists the comparison stars used in the photoelectric magnitude measurements.

All measurements were corrected for both atmospheric extinction and transformation. Transformation coefficients were computed using the two-star method outlined in Hall and Genet (1988, 200). The transformation coefficients for Frank's equipment are:  $\epsilon_V = -0.038$  and  $\epsilon_I = -0.16$ .

**Table 1: Characteristics of the 2000 apparitions of Uranus, Neptune and Pluto.<sup>a</sup>**

Parameter	Uranus	Neptune	Pluto
First conjunction date	2000 Feb. 6	2000 Jan. 24	1999 Dec. 3
Opposition date	2000 Aug. 11	2000 July 27	2000 June 1
Angular diameter (opposition)	3" .7	2" .4	0" .1
Right Ascension	21 <sup>h</sup> 26 <sup>m</sup>	20 <sup>h</sup> 30 <sup>m</sup>	16 <sup>h</sup> 46 <sup>m</sup>
Declination	-15° 52 <sup>m</sup>	-18° 46 <sup>m</sup>	-10° 57 <sup>m</sup>
Second conjunction date	2001 Feb. 9	2001 Jan. 26	2000 Dec. 7

<sup>a</sup> Data is taken from the *Astronomical Almanac* ©1998, 1999, and 2000a

**Table 2: People who contributed remote planets observations in 2000.**

Observer and Location	Type of Observation *
Phil Barker; Christchurch, New Zealand	V
Ray Berg; Crown Point, IN, USA	C, V, VP
Norman Boisclair; South Glens Falls, NY, USA	C, V
Brian Cudnik; Weimar, TX, USA	C, V, VP
Robert English; Franklin, TN, USA	V
Robin Gray; Winnemucca, NV, USA	C, V, VP
Walter Haas; Las Cruces, NM, USA	H, VP
Rik Hill; near Tucson, AZ, USA	VP
Raffaello Lena and others; Italy	CCD, V
Brian Loader; Christchurch, New Zealand	PP
Frank Melillo; Holtsville, NY, USA	CCD, CCD-P, PP, S, M
Katie Moore; Williamson, GA, USA	PP
Richard Schmude, Jr.; Barnesville, GA, USA	VP, PP
Tõfol Tobal and others; Spain	V
Roger Venable; Augusta, GA, USA	V
John Westfall; Antioch, CA, USA	PP
*C = Color; CCD = CCD image, CCD-P = CCD photometry, H = Historical data, M = methane band data, PP = photoelectric photometry, S = Spectra, V = visual observations or drawing, VP = eyeball magnitude	

John reports a value of  $\epsilon_I = -0.129 \pm 0.017$ . Loader's transformation coefficients are reported in (Schmude, 2000b, 159). During May/June, the author used a 0.51 meter Newtonian telescope along with a photometer having transformation coefficients of:  $\epsilon_B = 0.037$ ,  $\epsilon_V = -0.065$ ,  $\epsilon_R = -0.024$  and  $\epsilon_I = -0.135$ . In September, the author and Ms. Moore used a 0.09 meter Maksutov telescope and photometer having values of:  $\epsilon_B = 0.092$ ,  $\epsilon_V = -0.051$ ,  $\epsilon_R = -0.021$  and  $\epsilon_I = -0.095$ .

All photoelectric magnitude measurements of Uranus are listed in Table 4 while those of Neptune are listed in Table 5. In most cases, a measurement consisted of the sequence CPCPCPC where C is the comparison star reading and P is the Uranus (or Neptune) reading. Except in rare cases, each measurement is the average of between 3 and 5 P measurements. The selected normalized magnitudes are listed in Table 6.

The V-filter measurements of Loader, Melillo, Moore and Schmude correspond to normalized magnitudes of -7.12, -7.14, -7.15 and -7.15, which are all consistent

with one another. The infrared magnitudes of Melillo and Westfall correspond to normalized magnitudes of: -5.88 and -5.61; the difference between these two values is 0.27 magnitudes, which is too large. As a result of this discrepancy, I would like to recommend four things for all future photoelectric magnitude measurements: 1) always use a check star in making measurements, 2) take several sets of sky, comparison star, target, and check star measurements, 3) use a comparison star that has a B-V value close to that of the target object and 4) use a dew shield when making measurements.

The V(1,0) values of Uranus and Neptune over the last 10 years are plotted in Figure 1. Small color corrections were made to the 1991-1995 and 1997 data since these corrections were not made in the original papers. The data were fitted to a linear least squares routine and it was found that Uranus became 0.0020 magnitudes per year dimmer during the 1990s while Neptune became 0.0112 magnitudes per year brighter during that same time period. Most of the dimming for Uranus, however, took place between 1991 and 1996; during these years, Uranus became dimmer at an average annual rate of 0.0048 magnitudes. Lockwood and Thompson (1999, 3) carried out an excellent study of Uranus using the Strömgren b (4720 Å) and y (5510 Å) filters where the peak wavelength is given in parentheses. (The symbol Å means angstrom;  $1.0 \text{ Å} = 1.0 \times 10^{-10}$  meter). Lockwood and Thompson's study extends from 1972 through 1996. Their y-filter data shows Uranus getting 0.1 magnitudes brighter between 1972 and 1985 and getting 0.06 magnitudes dimmer between 1989 and 1996. The average dimming rate between 1989 and 1996 is around 0.008 magnitudes per year, which is larger than the 0.0048 magnitude/year rate measured by ALPO members. The higher dimming rate measured by Lockwood and Thompson may be due to the y-filter being almost centered on a methane absorption band; see Figure 2. If methane absorption had increased, then it would explain the extra dimming of Uranus in the y-filter. The Johnson V filter used by ALPO members, however, would be less affected by increased methane absorption since it covers a large wavelength range; see Figure 2.

During the 1990s, Neptune was brightest around 1999; this is 1 to 2 years from solar maximum. This result is different from the trend observed in the 1970s and 1980s, where Neptune was dimmest at solar maximum (Lockwood and Thompson, 1986, 1543). Hammel and Lockwood (1997, 466) report that a dark belt became less distinct in the mid 1990s, which may explain the 0.04 magnitude brightness increase at a wavelength of 5510 Å which they observed between 1989 and 1994. The brightening trend of Neptune shown in Figure 1 is also consistent with the results of Lockwood and Hammel (2000, 1009).

**Table 3: Summary of the comparison stars used in the photoelectric and CCD magnitude measurements made in 2000.**

Star Name	Right Ascension <sup>a</sup>	Declination <sup>a</sup>	Magnitude			
			B	V	R	I
Gamma-Cap	21 <sup>h</sup> 40.1 <sup>m</sup>	-16° 40 <sup>m</sup>	---	---	---	3.22 <sup>b</sup>
45-Cap	21 <sup>h</sup> 44.0 <sup>m</sup>	-14° 45 <sup>m</sup>	---	5.96 <sup>c,d</sup>	---	---
Theta-Cap	21 <sup>h</sup> 05.9 <sup>m</sup>	-17° 14 <sup>m</sup>	4.08 <sup>b</sup>	4.07 <sup>b</sup>	4.06 <sup>b</sup>	---
Rho-Cap	20 <sup>h</sup> 28.9 <sup>m</sup>	-17° 49 <sup>m</sup>	5.18	4.80	---	---
HD 195859	20 <sup>h</sup> 34.5 <sup>m</sup>	-19° 24 <sup>m</sup>	---	7.82 <sup>c,e</sup>	---	---
Star a	16 <sup>h</sup> 40.80 <sup>m</sup>	-10° 57.4 <sup>m</sup>	---	~12.5 <sup>f</sup>	---	---
Star b	16 <sup>h</sup> 42.95 <sup>m</sup>	-10° 57.4 <sup>m</sup>	---	~13.9 <sup>f</sup>	---	---
<sup>a</sup> All positions are from Hirshfeld, Sinnott and Ochsenbein (1991).						
<sup>b</sup> Magnitudes are from Iriarte et al. (1965).						
<sup>c</sup> Magnitudes are from the Tycho Catalog						
<sup>d</sup> A value of B-V = 0.21 was used for the transformation corrections; this B-V value is from Hirshfeld, Sinnott and Ochsenbein (1991).						
<sup>e</sup> A value of B-V = 0.4 was used for the transformation corrections; this B-V value is from Hirshfeld, Sinnott and Ochsenbein (1991).						
<sup>f</sup> Based on an average Pluto magnitude of 13.7 and unfiltered CCD images.						

### “Eyeball” Magnitude Estimates

Six individuals made 77 and 60 eyeball magnitude estimates of Uranus and Neptune respectively in 2000. Essentially, these people compared the brightness of these two planets to nearby comparison stars with only a telescope or binoculars. The average normalized magnitudes are:  $-7.1 \pm 0.01$  and  $-6.9 \pm 0.01$  for Uranus and Neptune respectively. Only random errors are included in the uncertainty for the normalized magnitudes. Hollis (2000, 124) reports a mean brightness magnitude for Uranus of  $+5.59 \pm 0.171$ , which is equivalent to  $V_{vis}(1,0) = -7.12$ ; this is close to the value selected for the 2000 apparition. Hollis (1999, 278) also reports a mean brightness of  $7.76 \pm 0.19$  for Neptune between 1989 and 1997 which corresponds to a mean normalized magnitude of  $-6.95$ .

### Appearance of Uranus and Neptune

Phil Barker and Robin Gray suspected a bright area on Uranus near the limb on June 8 and Oct. 17 respectively. A similar spot was observed by myself 12 years earlier and was independently confirmed by others (Schmude, 1988, 87). Limb darkening was noted in about half of the Uranus and Neptune observations. Brian Cudnik suspected a bright area near Neptune’s north pole that was bordered by a darker band. No other disc irregulari-

ties were reported on Neptune except for limb darkening. Twelve color estimates were reported for Uranus and of these, there were an equal number of blue and green hues reported. I have tabulated the color estimates of Uranus based on the colors of blue and green; the results are: blue (3), green (3), blue-green (3), blue to blue green (2) and yellow (1); the number of observations for each color is in parentheses. It is concluded that Uranus had an equal amount of blue and green color in 2000. The color estimates for Neptune are: blue (4) and blue-green (1) and so it is concluded that Neptune displayed a bluish hue with little or no green in 2000. These selected colors are also

in agreement with observations made by European observers.

Roger Venable reports that he got better views of Uranus using the traditional one-eye view compared to using a binocular viewer. This is probably due to the binocular viewer splitting the light in half and giving each eye half of the light that it would receive through just a single eyepiece.

Frank Melillo used a 0.20 meter telescope along with a CCD camera on Sept. 1, 2000 (3:30 U.T.) to image Uranus; the image scale was 1 pixel = 0.5 arc-seconds. Frank reports no cloud features on Uranus, but he does report a pronounced limb darkening of the planet.

Frank Melillo used a methane band filter with a peak transmission of light having a wavelength of 8900 Å along with a 0.20 meter telescope and a CCD camera to image Uranus. The methane band filter was purchased from Andover Corporation of Salem, New Hampshire and is 0.0070 meters thick. The images made with this filter are called “methane band images” because 8900 Å is a wavelength of light that methane absorbs; methane band images of Uranus and Neptune are more likely to show albedo features than integrated light images. A total of 16 images were made on Aug. 22, 23, Sept. 7, 8 and 22, and from these, Frank detected no change in brightness and concludes that any change would be less

**Table 4: Measured and normalized magnitudes of Uranus made in 2000 based on photoelectric measurements.**

Date – U.T.	Filter	Magnitude		Comparison Star
		Measured (+)	Normalized (-)	
May 7.386	V	5.85	7.10	Gamma-Cap
May 16.763	V	5.87	7.12	45-Cap
June 7.316	V	5.82	7.13	Gamma-Cap
June 7.351	V	5.79	7.16	"
June 7.357	V	5.77	7.18	"
Aug. 6.229	V	5.75	7.13 <sup>a</sup>	Theta-Cap
Aug. 6.240	I	7.03	5.86 <sup>a</sup>	"
Aug. 17.177	V	5.75	7.13 <sup>a</sup>	"
Aug. 17.188	I	7.01	5.88 <sup>a</sup>	"
Aug. 21.167	V	5.75	7.14 <sup>a</sup>	"
Aug. 21.177	I	6.98	5.90 <sup>a</sup>	"
Sept. 5.232	I	7.27	5.62	Gamma-Cap
Sept. 6.266	I	7.27	5.63	"
Sept. 7.258	I	7.27	5.62	"
Sept. 8.267	I	7.23	5.67	"
Sept. 18.104	V	5.77	7.14 <sup>a</sup>	Theta-Cap
Sept. 18.115	I	7.05	5.86 <sup>a</sup>	"
Sept. 19.281	I	7.30	5.62	Gamma-Cap
Sept. 20.230	I	7.32	5.59	Gamma-Cap
Sept. 21.252	I	7.32	5.60	"
Sept. 24.235	I	7.33	5.59	"
Sept. 25.261	I	7.30	5.62	"
Sept. 27.221	I	7.31	5.62	"
Sept. 28.107	B	6.36	6.56	Theta-Cap
Sept. 28.078	V	5.78	7.14	"
Sept. 28.134	R	6.10	6.83	"
Sept. 29.076	R	6.12	6.80	"
Sept. 29.095	R	6.13	6.80	"
Sept. 29.114	R	6.11	6.81	"
Sept. 29.134	R	6.04	6.88	"
Sept. 29.153	R	6.10	6.83	"
Sept. 29.176	R	6.11	6.82	"
Sept. 29.197	R	6.09	6.84	"
Sept. 29.209	I	7.35	5.58	"
Sept. 29.215	R	6.14	6.78	"
Oct. 2.193	I	7.31	5.62	Gamma-Cap
Oct. 4.220	I	7.31	5.62	"
Oct. 7.236	I	7.39	5.55	Gamma-Cap
Oct. 17.197	I	7.35	5.61	"
Oct. 19.220	I	7.43	5.53	"
Oct. 22.144	I	7.33	5.63	"
Oct. 29.025	V	5.78	7.19	Theta-Cap
Oct. 29.049	V	5.82	7.15	"
Nov. 1.155	I	7.33	5.65	Gamma-Cap
Nov. 17.116	I	7.38	5.63	"

<sup>a</sup> These measurements are based on just one set of measurements and are given a ¼ weight.

than 0.1 to 0.2 magnitudes. Unfortunately, Frank was unable to make quantitative brightness estimates of Uranus due to a low signal-to-noise ratio.

## Spectroscopy

Melillo imaged the spectra of both Uranus and Neptune; the dates and times of the Neptune spectra are: August 17, 2000 (4:15 U.T.) and Aug. 21, 2000 (4:15 U.T.). One of his spectra is shown in Figure 2. Essentially, Melillo used a spectroscope sold by Rainbow Optics and imaged the spectra with a CCD camera. He used Starlight Express software for Uranus to first stretch the spectrum and then take an intensity row in the spectrum, which was converted into the intensity-versus-wavelength graph in Figure 2. The procedure was the same for Neptune, except that Maxlm DL software was used. The sharp drops in intensity in Figure 2 are called absorption bands. The absorption bands near 5400, 6200, 7200, 7950, 8900 and 9900 Å are caused by the methane in the atmosphere of Uranus absorbing light at these wavelengths. Other absorption bands near 5750, 6670, 6970 and 8540 Å are present in the Uranus spectrum while additional absorption bands at 5740, 6640, 6890 and 7850 Å are present in the Neptune spectrum. The intensity of light at 4500 Å divided by that at 5400 Å was measured from the spectra of both planets; the resulting quotients are: 0.7 for Uranus and 0.5 for Neptune. These results may be due to differences in the atmospheric transparency on the dates that the spectra were imaged, or the spectra may show that the reflectivity of Neptune drops off more steeply at shorter wavelengths.

The ratio of the depth of the methane absorption bands near 5400, 6200 and 7200 Å to the peak intensity were evaluated for 1999 and 2000. The bands near 5400 and 6200 Å were deeper in 1999 than in 2000, but the reverse was the case for the 7200 Å band, and so no firm conclusions can be made regarding changes in the methane absorption bands between 1999 and 2000. It is important for members to continue to record the spectra of Uranus and Neptune in the future so that the methane absorption bands can be monitored.

## Titania, Oberon and Triton

Melillo recorded two unfiltered CCD images of Uranus, Titania and Oberon on

**Table 5: Measured and normalized magnitudes of Neptune made in 2000 based on photoelectric measurements**

Date – U.T.	Filter	Magnitude		Comparison Star
		Measured (+)	Normalized (-)	
May 1.382	V	7.74	7.04	Rho-Cap
May 16.739	V	7.79	6.97	HD 195859
June 7.292	V	7.74	7.00	Rho-Cap
June 7.334	V	7.79	6.95	“
June 7.375	V	7.78	6.96	“

Aug. 6 (5:45 U.T.) under excellent seeing conditions. Titania was 0.22 and 0.27 magnitudes brighter than Oberon in two different frames. These results are intermediate between those in 1998 and 1999. For 1998-2000, Titania is, on average,  $0.21 \pm 0.04$  magnitudes brighter than Oberon which is consistent with the literature value (*Astronomical Almanac*, 2000b). Melillo also recorded images of Neptune and Triton on Aug. 17 (4:10 U.T.) and Aug. 21 (4:04 U.T.). Melillo reports that Triton may be fainter at southern elongation than at western elongation; unfortunately, the interference from Neptune was too great to do precise photometry of Triton by itself.

## Pluto

Melillo made a series of CCD images of the Pluto system, hereafter referred to as “Pluto”, along with two comparison stars (“stars a” and “b” in Table 3) between July 2 and 9. He essentially imaged Pluto along with two comparison stars using two 30-second exposures, which were combined into one image. The relative brightness of Pluto and star a was measured for each date; furthermore, the relative brightness between stars a and b was measured and it was found that the magnitude differences between these stars remained within 0.06 magnitudes of the average difference. The longitude of the sub-Earth point on Pluto (*Astronomical Almanac*, 1999, E78) was also computed for each of Frank’s measurements; the results are shown in Figure 3. The 1998 Pluto data summarized elsewhere (Schmude, 2000a, 17) were also plotted as a function of the longitude of the

sub-Earth point on Pluto (*Astronomical Almanac*, 1997, E78) and this is also shown in Figure 3. Different comparison stars were used in 1998 and 2000, and so one cannot combine the data into one graph; nevertheless, one can see an overall agreement between the two graphs. The two brightness minima recorded by Melillo were compared to a brightness minimum measured by Binzel and Mulholland (1984, 1760) on Julian date 2,445,434.763. Melillo’s two brightness minima were measured on June 2.16, 1998 (Julian date = 2,450,966.66) and July 7.16, 2000 (Julian date = 2,451,732.66). The difference between Melillo’s two times is 766.0 days which equals 119.93 Pluto rotations while the difference between Frank’s 2000 measurement and Binzel and Mulholland’s measurement is 6297.897 days, which equals 986.02 Pluto rotations; a rotational period of 6.3872 days for Pluto was used (*Astronomical Almanac*, 2000b, E88). These results confirm the stability of the bright and dark areas on Pluto. I conclude that the Pluto system is brightest when the sub-Earth longitude is near  $130^\circ$  and is dimmest near  $280^\circ$ .

The difference between maximum and minimum brightness as Pluto rotates has changed from 0.14 magnitudes in 1954 to 0.28 magnitudes in 1982 (Stern, Trafton and Gladstone, 1988, 486). In 1998-2000, the variation was near 0.4 magnitudes. Stern, Trafton and Gladstone (1988, 485) predict that the variation in Pluto’s rotation curve will begin to decline between 1996 and 2006. It is important for ALPO members to continue to monitor the brightness of Pluto as it rotates. Another prediction by Stern and co-workers was that Pluto’s geometric albedo will begin to rise after 1989. (The year 1989 was the date of perihelion passage.)

**Table 6: Selected normalized magnitudes for Uranus and Neptune in 2000.**

Planet	B(1,0)	V(1,0)	R(1,0)	I(1,0)
Uranus	-6.56	-7.14±0.01	-6.82±0.01	-5.62±0.05
Neptune	---	-6.98±0.02	---	---




Figure 1: Plots of the normalized magnitudes of Uranus and Neptune versus the year from 1991 to 2000.




Figure 2: The spectrum of Uranus recorded by Frank Melillo on Aug. 27, 2000 at 5:30 U.T. Also shown are the effective wavelengths and bandpasses for the Johnson V filter (designated as V) and the Strömrgren y filter (designated as y).




Figure 3: Plots of the difference in magnitude between the Pluto system and selected comparison stars in 1998 and 2000 versus the longitude of the sub-Earth point on Pluto.

## References

- Astronomical Almanac for the year 1998 (1997)* Washington D.C.: U.S. Govt. Printing Office.
- Astronomical Almanac for the year 1999 (1998)* Washington D.C.: U.S. Govt. Printing Office.
- Astronomical Almanac for the year 2000 (1999)* Washington D.C.: U.S. Govt. Printing Office.
- Astronomical Almanac for the year 2001 (2000a)* Washington D.C.: U.S. Govt. Printing Office.
- Astronomical Almanac for the year 2002 (2000b)* Washington D.C.: U.S. Govt. Printing Office.
- Binzel, R. P. and Mulholland, J. D. (1984) "Photometry of Pluto During the 1983 Opposition: A New Determination of the Phase Coefficient." *The Astronomical Journal*, Vol. 89, No. 11, pp. 1759-1761.
- Hall, D. S. and Genet, R. M. (1988) *Photoelectric Photometry of Variable Stars*, Second Edition, Richmond VA: Willmann-Bell, Inc.
- Hammel, H. B. and Lockwood, G. W. (1997) "Atmospheric Structure of Neptune in 1994, 1995 and 1996: HST Imaging at Multiple Wavelengths." *Icarus*, Vol. 129, pp. 466-481.
- Hirshfeld, A., Sinnott, R. W. and Ochsenein, F. (1991) *Sky Catalog 2000.0*, Vol. 1, Cambridge MA: Sky Publishing Corp.
- Hollis, A. J. (1999) "Neptune 1989-1997." *Journal of the British Astronomical Assn.*, Vol. 109, No. 5, pp. 278-280.
- Hollis, A. J. (2000) "Uranus 1987-1997." *Journal of the British Astronomical Assn.*, Vol. 110, No. 3, pp. 124-126.
- Iriarte, B., Johnson, H. L., Mitchell, R. I. And Wisniewski, W. K. (1965) "Five-Color Photometry of Bright Stars." *Sky & Telescope*, Vol. 30, No. 1 (July) pp. 21-31.
- Lockwood, G. W. and Hammel, H. B. (2000) "Neptune's Increasing Brightness Since 1989: Inferences from Photometry, Voyager and HST Images, and Recent Keck Adaptive Optics Images." *Bulletin of the American Astronomical Society*, Vol. 32, p.1009.
- Lockwood, G. W. and Thompson, D. T. (1986) "Long-Term Brightness Variations of Neptune and the Solar Cycle Modulation of its Albedo." *Science*, Vol. 234, pp. 1543-1545.
- Lockwood, G. W. and Thompson, D. T. (1999) "Photometric Variability of Uranus, 1972-1996." *Icarus*, Vol. 137, pp. 2-12.
- Optec Inc. (1997) *Model SSP-3 Solid-State Stellar Photometer Technical Manual for Theory of Operation and Operating Procedures*, revision three, Lowell, MI: Optec Inc.
- Roddier, F., Roddier, C., Graves, J. E., Northcott, M. J. and Owen, T. (1998) "Neptune's Cloud Structure and Activity: Ground-Based Monitoring with Adaptive Optics." *Icarus*, Vol. 136, pp. 168-172.
- Schmude, R. W. Jr. (1988) "A View of Uranus and Neptune." *National Newsletter of the Royal Astronomical Society of Canada*, Vol. 82, pp. 86-87.
- Schmude, R. W. Jr. (1992) "The 1991 Apparition of Uranus." *Journal of the Association of Lunar & Planetary Observers*, Vol. 36, No. 1 (March) pp. 20-22.
- Schmude, R. W. Jr. (2000a) "Observations of the Remote Planets in 1998." *Journal of the Association of Lunar & Planetary Observers*, Vol. 42, No. 1 (January) pp. 13-17.
- Schmude, R. W. Jr. (2000b) "Observations of the Remote Planets in 1999." *Journal of the Association of Lunar & Planetary Observers*, Vol. 42, No. 4 (October) pp. 158-162.
- Sromovsky, L. A., Spencer, J. R., Baines, K. H. and Fry, P. M. (2000) "Ground-Based Observations of Cloud Features on Uranus." *Icarus*, Vol. 146, pp. 307-311.
- Stern, S. A., Trafton, L. M. and Gladstone, G. R. (1988) "Why is Pluto Bright? Implications of the Albedo and Lightcurve Behavior of Pluto." *Icarus*, Vol. 75, pp. 485-498.
- Talcott, R. (2001) "Close-up on Distant Giants." *Astronomy*, Vol. 29, No. 3 (March) pp. 30-32.
- Wasserman, L. H., Olkin, C. B. and Franz, O. G. (2000) "HST Measurement of the Charon/Pluto Mass Ratio." *Bulletin of the American Astronomical Society*, Vol. 32, p. 1082.
- Young, E. F., Binzel, R. P. and Crane, K., (2001) "A Two-Color Map of Pluto's Sub-Charon Hemisphere." *The Astronomical Journal*, Vol. 121, pp. 552-561.

**ALPO Feature:  
Index to Volume 42 of the “Strolling Astronomer”  
Journal of the Assn. of Lunar & Planetary Observers**

**By Michael Mattei**

**Publication Data**

- Issue Number: 1, January, 2000.....pp. 1-48
- 2, April, 2000.....pp. 49-96
- 3, July, 2000.....pp. 97-148
- 4, October, 2000....pp. 149-196

(Abbreviations: I. = Issue, IBC = Inside Back Cover,  
OFC = Outside Front Cover)

**Author Index**

- Baum, Richard  
The Enigmatic Light of Venus: An Overview  
..... 118-125
- Beish, Jeffrey D.  
Drawing Mars ..... 36-38
- Benton, Julius L.  
Observations of Saturn During the 1996-97  
Apparition ..... 1-12
- Observations of Saturn During the 1997-98  
Apparition ..... 106-117
- Observations of Venus During the 1996-97  
Western (Morning) Apparition ..... 49-57
- Observations of Venus During the 1997-98  
Eastern (Evening) Apparition ..... 149-157
- Some Recollections of Youth, the ALPO,  
and Observations of Saturn..... 30-35
- Budosh, Carole J.  
Charles Morgan Cyrus ..... 35
- Dobbins, Thomas, Sheehan, William  
Project “Delta Luna”: A Proposal to Search for  
Impact Features on the Moon of Recent Origin  
..... 172-174
- Favero, Giancarlo  
The Linne’ Controversy; or On the Shape of Lunar  
Craterlets Seen Under Different Solar Altitudes  
..... 186-190
- Favero, Giancarlo, Lena, Raffaello, Mengoli, Giorgio,  
Cipolat, Alessandro, Gualdoni, Piermario;  
The Nature of the Hook-like Shadow

Observed by Wilkins and Moore on Plato’s Floor in  
1952 ..... 126-132

- Garrett, Lawrence  
ALPO Minor Planets Section Report and Review for  
1999: Part I Review of the Minor Planets Section  
..... 133-136
- ALPO Minor Planets Section Report and Review for  
1999: Part II Magnitude Alert Project: Suggested  
Revised H-Values of Selected Asteroids for 1999  
..... 136-137
- Gingrich, Mark  
Illuminated Extent: A Telltale Measure of an Inferior  
Planet’s Disk ..... 18-22
- Haas, Walter H.  
A Visual Observational Study of the Extensions of  
the Cusps of Venus, 1937-1998 ..... 58-66
- Jamieson, Harry D.  
The New ALPO Journal and What it Means to You  
..... 138-139
- Kronk, Gary  
Aims of the Comets Section ..... 88-89
- Machholz, Don  
The Coming Apparition of Comet Linear  
(C/199 S4) ..... 23-25
- Comet Hyakutake, C/1996 B2..... 97-105
- Mattei, Michael  
Index to Volume 42 of the Journal of the Association  
of the Lunar & Planetary Observers ..... 41-43
- Melillo Frank J.  
CCD Methane Band Observations of Titan in 1999  
..... 175-176
- Minton, R.B.  
Thrift Store Computers versus Supercomputers 28-29
- Pilcher, Frederick  
ALPO Minor Planets Section Report and Review for  
1999: Part III General Report of Position Observa-  
tions by the ALPO Minor Planets Section .. 163-171
- Poshedly, Ken  
Memorable Moments at the Peach State Star Gaze  
..... 26-27
- About Antoni’n Rukl ..... 27

## The Strolling Astronomer

Schmude, Richard

- Observations of the Remote Planets in 1998... 13-17
- Observations of the Remote Planets in 1999  
..... 158-162
- The ALPO Historical Section..... 29
- Wideband Photometry of Jupiter: 1999-2000..67-72

Salimbeni, P.G., Lena, R., Mengoli, E., Douglas, E.,  
Santacana; Geologic Lunar Research Group (GLR)  
The Domes of the Rima Birt Region ..... 83-87

Westfall, Elizabeth

- Minutes of the Board of Directors Meeting, ALPO,  
Ventura, California, July 19-22, 2000..... 140-141

Westfall, John E.

- Whole-Disk Photometry of Jupiter:1991-1994  
..... 72-73
- Galilean Satellite Eclipse Timings:  
The 1994/95 Apparition..... 74-82

Williams, Thomas R.

- A Brief History of the ALPO Meteors  
Project..... 177-185

### Subject Index

ALPO

- Board of Directors ..... 47, 95, 146, 194
- Board and Staff Internet Directory  
..... IBC, Nos. 1, 2, 3 4
- Paper Session at Astrocon, July 20, 2000..... 90
- Staff.....48-IBC, 96-IBC, 146-148, 194-196
- The Association of Lunar and Planetary Observers  
..... 44,92

ALPO Announcements, Section Changes, Other

ALPO News

- Address Changes ..... 143
- ALPO Web Master..... 143
- Brad Smith Honored ..... 191
- Convention in 2001..... 92
- E-Mail Address Changes ..... 190
- New Appointments for Expanded Publications  
Section ..... 142
- New Comets Coordinator ..... 44
- New Instrument Section Staff..... 142
- New Meteorite Section ..... 142
- New Minor Planet Section Staff ..... 142
- New Name and Coordinator for Youth programs  
Section ..... 142
- Observing and Understanding Uranus, Neptune and  
Pluto ..... 191
- Recent Board Decisions..... 92
- Reminder for Authors ..... 191
- Staff and Section Changes ..... 92
- Welcome Donations ..... 143

Other Amateur and Professional Announcements

- Consolidated Lunar Atlas On Line..... 191
- Deadline for NSF Planetary Science Proposals.... 93
- Double Star Observer ..... 45

Roster of Upcoming Meetings

- July 9-12, 2000: Catastrophic Events and Mass  
Extinctions: Impacts and Beyond ..... 45
- July 9-14, 2000: International Planetarium Society  
Conference..... 45
- July 19-22, 2000: Astrocon 2000 ..... 45
- August 7-18, 2000: International Astronomical Union  
General Assembly and Associated Symposia ..... 45
- August 21-25, 2000: Second International Confer-  
ence on Mars Polar Science and Exploration ..... 45
- August 28-September 1, 2000: 63rd Annual Meeting  
of the Meteoritical Society ..... 45
- September 22-23, 2000: Nightfall ..... 93
- September 22-24, 2000: MERAL/VAAS Convention  
and Star Party 2000 ..... 93
- September 28-October 1, 2000: Enchanted Skies Star  
Party ..... 93
- October 14-15, 2000: Solar Eclipse Conference 2000  
..... 45, 93
- October 23-27, 2000: 32nd Division for Planetary  
Science Meeting ..... 45, 93
- November 8-10, 2000: Conference on the Earth-  
Moon Relationship ..... 93
- November 13-16, 2000: Annual Meeting of the Geo-  
logical Society of America ..... 93
- December 5-7, 2000: ISCO 2000, International Con-  
ference on Space Optics ..... 93, 143
- January 4-8, 2001: Small-Telescope Astronomy on  
Global Scales ..... 93, 143
- January 7-11, 2001: 197th Meeting of the American  
Astronomical Society ..... 143
- March 19-24, 2001: International Planetarium and  
Astronomy Conference..... 191
- March 26-28, 2001: Fourth Annual Raytheon Science  
Data Centers Symposium ..... 191
- April 2-7, 2001 Brown Dwarfs and Planet, 31st Saas-  
Fee Advanced Course..... 191
- June 11-15, 2001: First Eddington Workshop .... 191
- June 18-20, 2001: Workshop on the Evolving Sun  
and Its Influence on Planetary Enviroments..... 191
- June 25-27, 2001: Conference on Jupiter-Planet,  
Satellites and Magnetosphere ..... 93, 143, 191
- July 24-28, 2001: ALCON 2001 ..... 93, 143, 191

ALPO Convention

- Address and Address Changes ..... 44, 92
- Awards for the Year 2000 ..... 93
- Generous Members..... 45
- Minor Planet Elmerreese ..... 44-45

Book Reviews (Edited by Jose Olivarez)

- Atlas of the Lunar Terminator ..... 185

## The Strolling Astronomer

Comets	
Aims of the Section .....	88-89
Coming Apparition of Comet Linear (C/199 S4)	
.....	23-25
Hyakutake, C/1996 B2.....	97-105
History	
Fifty Years Ago.....	40,66
Historical Section.....	29
Index	
Index to Volume 41 of the J.ALPO.....	41-43
In Memoriam	
Charles Morgan Cyrus .....	35
Instruments	
Thrift Store Computers versus Supercomputers.....	28-29
Journal of the ALPO	
New Journal and What it Means to You .....	138
Jupiter (See also under "satellites")	
Whole-Disk Photometry of Jupiter:1991-1994	
.....	72-73
Wideband Photometry of Jupiter: 1999-2000..	67-72
Mars	
Drawing Mars .....	36-38
Meetings	
The ALPO Meets on the Shores of the Pacific:	
Astrocon 2000.....	89-91
Meteors	
Brief History of the ALPO Lunar Meteors	
Project .....	177-185
Minor Planets	
Report and Review for 1999: Part I Review of the	
Minor Planets Section .....	133-136
Report and Review for 1999: Part II Magnitude Alert	
Project: Suggested Revised H-Values	
of Selected Asteroids for 1999 .....	136-137
Report and Review for 1999: Part III - General	
Report of Position Observations by the Minor	
Planets Section.....	163-171
Moon; General	
Project "Delta Luna": A Proposal to Search for	
Impact Features of Recent Origin.....	172-174
Something Does Happen on the Moon .....	125
Moon; Craters, Features, and Regions	
Domes of the Rima Birt Region .....	83-87
Linne' Controversy; or On the Shape of Lunar Crater-	
lets Seen Under Different Solar Altitudes ...	186-190
Nature of the Hook-like Shadow Observed by	
Wilkins and Moore on Plato's Floor in 1952	126-132
New Books Received (Comments by Jose Olivarez)	
A Sky Watcher's Year .....	39
Meteorites and Their Parent Planets, Second Edition	
.....	40
The Deep Sky, An Introduction.....	39
The Monthly Sky Guide, 5th Edition .....	39
Observations	
Illuminated Extent: A Telltale Measure of an Inferior	
Planet's Disk .....	18-22
Other Publications of the ALPO	
An Introductory Bibliography for Solar System	
Observers .....	46, 94, 144, 192
Membership Directory.....	46, 94, 144, 193
Back Issues of J.ALPO.....	46, 94, 144, 193
Publications of the Sections of the ALPO	
Computing .....	47, 95,146,194
Jupiter .....	47, 95, 145, 194
Lunar and Planetary Training Program	
.....	47, 95, 145, 146, 193
Lunar.....	47, 95, 145, 193
Mars.....	47, 95, 145, 194
Meteors .....	47, 95, 146, 194
Minor Planets.....	47, 95, 146, 194
Saturn.....	47, 95, 145, 194
Venus .....	47, 95, 145, 193- 194
Publications, ALPO	
ALPO Monograph Series .....	46, 94, 143-144, 192
Remote Planets, Uranus, Neptune, Pluto	
Observations of the Remote Planets in 1998... 13-17	
Observations of the Remote Planets in 1999	
.....	158-162
Satellites	
CCD Methane Band Observations of Titan in 1999	
.....	175-176
Galilean Satellite Eclipse Timings: The 1994/95	
Apparition.....	74-82
Saturn	
Observations During the 1996-97 Apparition ... 1-12	
Observations During the 1997-98 Apparition.....	106-117
Venus	
Enigmatic Light of Venus: An Overview .... 118-125	
Observations During the 1996-97 Western (Morning)	
Apparition.....	49-57
Observations During the 1997-98 Eastern (Evening)	
Apparition.....	149-157
Visual Observational Study of the Extensions of the	
Cusps 1937-1998 .....	58-66

## **ALPO Resources**

**People, publications, etc. to help our members**

### **Board of Directors**

- Executive Director (Chair); Julius L. Benton, Jr., Associates in Astronomy, 305 Surrey Road, Savannah, GA 31410
- Associate Director; Donald C. Parker, 12911 Lerida Street, Coral Gables, FL 33156
- Founder/Director Emeritus; Walter H. Haas, 2225 Thomas Drive, Las Cruces, NM 88001
- Member of the Board; Richard Hill, Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721
- Member of the Board; Ken Poshedly, 1741 Bruckner Ct., Snellville, GA 30078-2784
- Member of the Board; Richard W. Schmude, Jr., 316 Forsyth St., #5, Barnesville, GA 30204
- Member of the Board, Secretary; Elizabeth W. Westfall, P.O. Box 2447, Antioch, CA 94531-2447
- Member of the Board; John E. Westfall, P.O. Box 2447, Antioch, CA 94531-2447
- Membership Secretary/Treasurer; Matthew Will, P.O. Box 13456, Springfield, IL 62791-3456

### **Publications Staff**

- Publisher (Acting); Ken Poshedly. Send ALL papers and articles to 1741 Bruckner Ct, Snellville, GA 30078-2784; e-mail poshedly@bellsouth.net

### **Science Editors**

- Dr. Klaus R Brasch, Executive Director, Office of Technology Transfer & Professor of Biology, California State University San Bernardino, 5500 University Parkway, San Bernardino, CA 92407
- Dr. Richard K Ulrich (Acting), Professor, Dept. Of Chemical Engineering, 3202 Bell Engineering Center, University of Arkansas, Fayetteville, AR 72701
- John E Westfall, 5061 Carbondale Way, Antioch, CA 94509

### **General Editors (Acting)**

- Special Projects; Robert A Garfinkle, F.R.A.S., 32924 Monrovia St., Union City, CA 94587-5433
- Magazine Markup and Assembly; Ken Poshedly, 1741 Bruckner Ct, Snellville, GA 30078-2784
- Special Projects; Roger J Venable, MD, 3405 Woodstone Pl., Augusta, GA 30909-1844

### **Book Review Editor**

- Jose Olivarez, 4705 SE 14th St., Ocala, FL 34471

### **Graphics (Acting)**

- John Sanford, P.O. Box 1000, Springville, CA 93265-1000

### **Staff Writers (Acting)**

- Eric Douglass, 10326 Tariaton Dr., Mechanicsville, VA 23116-5835
- James S Lamm, 9211 13th Avenue Circle NW, Bradenton, FL 34209
- Richard J Wessling, 5429 Overlook Dr., Milford, OH 45150-9651

### **Translators (Acting)**

- French Language Submissions; Richard J McKim, Cherry Tree Cottage, 16 Upper Main Street, Upper Benefield, Peterborough Pe8 5an United Kingdom
- Spanish Language Submissions; Guido E Santacana, Nuevo Laredo 1678, Venus Gardens, Rio Piedras, PR 00926

### **Lunar and Planetary Training Program**

- Coordinator; Timothy L. Robertson, 2010 Hillgate Way #L, Simi Valley, CA 93065

### **Solar Section**

- Coordinator, *Website, SolNet, Rotation Report*, handbook; Richard Hill, Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721
- Assistant Coordinator, Correspondence and New Observers; Gordon W. Garcia, 340 Illinois Boulevard, Hoffman Estates, IL 60194-3319
- Acting Assistant Coordinator; Tony Grigsby, 209 Hubbard Lane, Mt. Washington, KY 40047
- Acting Assistant Coordinator; Brad Timerson (use e-mail for correspondence, see Internet directory)

### **Mercury Section**

- Acting Coordinator; Frank J. Melillo, 14 Glen-Hollow Dr., E-#16, Holtsville, NY 11742

### **Venus Section**

- Coordinator; Julius L. Benton, Jr., Associates in Astronomy, 305 Surrey Road, Savannah, GA 31410

### **Mercury/Venus Transit Section**

- Coordinator; John E. Westfall, P.O. Box 2447, Antioch, CA 94531-2447

## ALPO Resources (Continued)

### **Lunar Section**

- Coordinator, *Selected Areas Program*; Julius L. Benton, Jr., Associates in Astronomy, 305 Surrey Road, Savannah, GA 31410
- Acting Coordinator, Lunar Meteoritic Impacts Search; Brian Cudnik, 1616 Hollister Road, #1616, Houston, TX 77040
- Coordinator, Lunar Transient Phenomena; David O. Darling, 416 W. Wilson St., Sun Prairie, WI 53590-2114
- Coordinator, Lunar Topographical Studies; William M. Dembowski, 219 Old Bedford Pike, Windber, PA 15963

### **Mars Section**

- Coordinator, *all observations, U.S. correspondence*; Daniel M. Troiani, 629 Verona Court, Schaumburg, IL 60193
- Assistant Coordinator; CCD Imaging, "Mars Project"; Donald C. Parker, 12911 Lerida Street, Coral Gables, FL 33156
- Assistant Coordinator; Daniel Joyce, 6203 N. Keeler Avenue, Chicago, IL 60648-5109
- Assistant Coordinator; Jeff D. Beish, 842 Virginia Ave., Lake Placid, FL 33852
- Acting Assistant Coordinator, Archivist; Deborah Hines, 2341 South 11th Avenue, Boardview, IL 60153

### **Minor Planets Section**

- Coordinator; Frederick Pilcher, Illinois College, Jacksonville, IL 62650.
- Acting Assistant Coordinator; Lawrence S. Garrett, 206 River Road, Fairfax, VT 05454
- Acting Assistant Coordinator; Richard Kowalski, 7630 Conrad Street, 761 Zephyrhills, FL 33544-2729
- Acting Scientific Advisor; Steve Larson, Lunar & Planetary Lab, University of Arizona, Tucson, AZ 85721

### **Jupiter Section**

- Acting Coordinator, Richard W. Schmude Jr., Gordon College, Division of Natural Sciences and Nursing, 419 College Drive, Barnesville, GA 30204
- Assistant Coordinator, Scientific Advisor; Sanjay Limaye, University of Wisconsin, Space Science and Engineering Center, Atmospheric Oceanic and Space Science Bldg. 1017, 1225 W. Dayton St., Madison, WI 53706
- Assistant Coordinator, Transit Timings; John McAnally, 2124 Wooded Acres, Waco, TX 76710

- Assistant Coordinator, Newsletter; Craig MacDougal, 2602 E. 98th Avenue, Tampa, FL 33612
- Acting Assistant Coordinator, CCD Images; Damian Peach, 237 Hillington Square, Greyfriars House, King's Lynn, Norfolk PE30 5HX, United Kingdom
- Assistant Coordinator, Scientific Advisor; Agustin Sanchez-Lavega, Departamento Fisica Aplicada I, E.T.S. Ingenieros, Universidad del Pais Vasco, Bilbao, Spain
- Assistant Coordinator, Galilean Satellites; John E. Westfall, P.O. Box 2447, Antioch, CA 94531-2447

### **Saturn Section**

- Coordinator, Julius L. Benton, Jr., Associates in Astronomy, 305 Surrey Road, Savannah, GA 31410

### **Remote Planets Section**

- Coordinator; Richard W. Schmude, Jr.; Gordon College, Division of Natural Sciences and Nursing, 419 College Drive, Barnesville, GA 30204

### **Comets Section**

- Acting Coordinator; Gary Kronk, 1117 Troy-O'Fallon Rd., Troy, IL 62294
- Assistant Coordinator; James V. Scotti, Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721

### **Meteors Section**

- Coordinator; Robert D. Lunsford, 161 Vance Street, Chula Vista, CA 91910
- Acting Assistant Coordinator; Robin Grey, P.O. Box 547, Winnemucca, NV 89446

### **Meteorites Section**

- Acting Coordinator; Dolores Hill, Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721

### **Computing Section**

- Coordinator; Mike W. McClure, 170 Prestwick Ct., Bowling Green, KY 42103

### **Youth Section**

- Coordinator; Timothy L. Robertson, 2010 Hillgate Way #L, Simi Valley, CA 93065

## ALPO Resources (continued)

### ALPO Board/Staff E-mail Directory

Beish, J.D. .... jeffbeish@htn.net  
Benton, J.L. .... jlbaina@msn.com  
Brasch, K.R. .... kbrasch@wiley.csusb.edu  
Baum, R. .... richardbaum@julianbaum.co.uk  
Cudnik, B. .... xflare@earthlink.net  
Darling, D.O. .... DOD121252@aol.com  
Dembowski, W.M. .... dembow@mail.third-wave.com  
Douglass, E. .... ejdftd@mindspring.com  
Garcia, G.W. .... gordg@megsinet.net  
Garfinkle, R.A. .... ragarf@earthlink.net  
Garrett, L.S. .... Lgasteroid@globalnetisp.net  
Grigsby, T. .... tony@alltel.net  
Haas, W.H. .... haasw@zianet.com  
Hill, D. .... dhill@lpl.arizona.edu  
Hill, R. .... rhill@lpl.arizona.edu  
Hines, D. .... cmptdevil@hotmail.com  
Joyce, D. .... djoyce@triton.cc.il.us  
Kowalski, R. .... bitnik@bitnik.com  
Kronk, G. .... kronk@amsmeteors.org  
Lamm, J.S. .... jlspacerox@aol.com  
Larson, J.S. .... slarson@lpl.arizona.edu  
Limaye, S. .... sanjayl@ssec.wisc.edu  
Lunsford, R.D. .... lunro.imo.usa@home.com  
MacDougal, C. .... macdouc@prodigy.net  
McAnally, J. .... CPAJohnM@aol.com  
McClure, M.W. .... mike.mcclure@kctcs.net  
McKim, R.J. .... rmckim5374@aol.com  
Melillo, F. .... FrankJ12@aol.com  
Olivarez, J. .... n/a  
Parker, D.C. .... park3232@bellsouth.net  
Peach, D. .... dpeach78@netscapeonline.co.uk  
Pilcher, F. .... pilcher@hilltop.ic.edu  
Poshedly, K. .... poshedly@bellsouth.net  
Reynolds, M. .... reynolds@cosc.org  
Robertson, T.J. .... cometman@cometman.net  
Sanchez-Lavega, A. .... wupsalaa@bic00.bi.ehu.es  
Sanford, J. .... johnsan@sosinet.net  
Santacana, G.E. .... laffitte@prtc.net  
Schmude, R.W. .... Schmude@Falcon.gdn.peachnet.edu  
Slaton, J.D. .... jd@justfun.org  
Timerson, B. .... bwtimer@eznet.net  
Troiani, D.M. .... dantroiani@earthlink.net  
Ulrich, R.K. .... rulrich@uark.edu  
Venable, R.J. .... rjvmd@knology.net  
Wessling, R.J. .... pinesop@aol.com  
Westfall, E.W. .... ewestfal@sfsu.edu  
Westfall, J.E. .... 73737.1102@compuserve.com  
Will, M. .... will008@attglobal.net

### Historical Section

- Coordinator; Richard Baum, 25 Whitchurch Rd., Chester, CH3 5QA, United Kingdom

### Instruments Section

- Acting Coordinator; R.B. Minton, 568 N. 1st Street, Raton, NM 87740
- Acting Assistant Coordinator; Richard J Wessling, 5429 Overlook Dr., Milford, OH 45150-9651

### Website

- Coordinator; Richard Hill, Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721
- Acting Assistant Coordinator; Jonathan D. Slaton, 1025 George, Alton, IL 62002-2561

### Eclipse Section

- Acting Coordinator; Michael D. Reynolds, Chabot Observatory and Science Center, 10902 Skyline Boulevard, Oakland, CA 94619

## ALPO Publications: The Monograph Series

ALPO monographs are publications that we believe will appeal to our members, but which are too lengthy for publication in *The Strolling Astronomer*. They should be ordered from *The Strolling Astronomer* editor (P.O. Box 2447, Antioch, CA 94531-2447 U.S.A.) for the prices indicated, which include postage. Checks should be in U.S. funds, payable to "ALPO".

- **Monograph Number 1. *Proceedings of the 43rd Convention of the Association of Lunar and Planetary Observers. Las Cruces, New Mexico, August 4-7, 1993.*** 77 pages. Price: \$12 for the United States, Canada, and Mexico; \$16 elsewhere.
- **Monograph Number 2. *Proceedings of the 44th Convention of the Association of Lunar and Planetary Observers. Greenville, South Carolina, June 15-18, 1994.*** 52 pages. Price: \$7.50 for the United States, Canada, and Mexico; \$11 elsewhere.
- **Monograph Number 3. *H.P. Wilkins 300-inch Moon Map.*** 3rd Edition (1951), reduced to 50 inches diameter; 25 sections, 4 special charts; also 14 selected areas at 219 inches to the lunar diameter. Price: \$28 for the United States, Canada, and Mexico; \$40 elsewhere.
- **Monograph Number 4. *Proceedings of the 45th Convention of the Association of Lunar and Planetary Observers. Wichita, Kansas, August 1-5, 1995.*** 127 pages. Price: \$17 for the United States, Canada, and Mexico; \$26 elsewhere.

## ALPO Resources (continued)

- **Monograph Number 5. *Astronomical and Physical Observations of the Axis of Rotation and the Topography of the Planet Mars. First Memoir; 1877-1878.*** By Giovanni Virginio Schiaparelli, translated by William Sheehan. 59 pages. Price: \$10 for the United States, Canada, and Mexico; \$15 elsewhere.
- **Monograph Number 6. *Proceedings of the 47th Convention of the Association of Lunar and Planetary Observers, Tucson, Arizona, October 19-21, 1996.*** 20 pages. Price \$3 for the United States, Canada, and Mexico; \$4 elsewhere.
- **Monograph Number 7. *Proceedings of the 48th Convention of the Association of Lunar and Planetary Observers. Las Cruces, New Mexico, June 25-29, 1997.*** 76 pages. Price: \$12 for the United States, Canada, and Mexico; \$16 elsewhere.
- **Monograph Number 8. *Proceedings of the 49th Convention of the Association of Lunar and Planetary Observers. Atlanta, Georgia, July 9-11, 1998.*** 122 pages. Price: \$17 for the United States, Canada, and Mexico; \$26 elsewhere.
- **Monograph Number 9. *Does Anything Ever Happen on the Moon?*** By Walter H. Haas. Reprint of 1942 article. 54 pages. Price: \$6 for the United States, Canada, and Mexico; \$8 elsewhere.
- **Monograph Number 10. *Observing and Understanding Uranus, Neptune and Pluto.*** By Richard W. Schmude, Jr. 31 pages. Price: \$4 for the United States, Canada, and Mexico; \$5 elsewhere.
- **Lunar (Dembowski): *The Lunar Observer*,** a monthly newsletter, available online at the ALPO website homepage, <http://www.lpl.arizona.edu/alpo/>. For hard copies, send a set of self-addressed stamped envelopes to Bill Dembowski (see address in listings section earlier in this section).
- **Lunar (Jamieson): *Lunar Observer's Tool Kit*,** price \$25, is a PC program consisting of several scheduling and observing aids. Some of the program's options include predicting when a lunar feature will be illuminated in a certain way, what features from a collection of features will be under a given range of illumination, physical ephemeris information, mountain height computation, coordinate conversion, and browsing of the software's included database of over 6,000 lunar features. This program is DOS-based, and intended for more advanced lunar observers. Available by e-mail only; contact [hjamieso@telocity.com](mailto:hjamieso@telocity.com).
- **Venus (Benton):** (1) *The ALPO Venus Observing Kit*, \$17.50. Includes introductory description of ALPO Venus observing programs for beginners, a full set of observing forms, and a copy of *The Venus Handbook*. (2) *Observing Forms Packet*, \$10. Includes observing forms to replace those provided in the observing kit described above. Specify *Venus Forms*. To order either numbers (1) or (2), send a check or money order payable to "Julius L. Benton, Jr." All foreign orders should include \$5 additional for postage and handling; p/h included in price for domestic orders. Shipment will be made in two to three weeks under normal circumstances. NOTE: Observers who wish to make copies of the observing forms may instead send a SASE for a copy of forms available for each program. Authorization to duplicate forms is given only for the purpose of recording and submitting observations to the ALPO Venus, Saturn, or lunar SAP sections. Observers should make copies using high-quality paper.

## ALPO Observing Section Publications

Order the following directly from the appropriate ALPO section coordinators; use the address in the listings pages which appeared earlier in this booklet unless another address is given.

- **Lunar and Planetary Training Program (Robertson): *The Novice Observers Handbook*** \$15. An introductory text to the training program. Includes directions for recording lunar and planetary observations, useful exercises for determining observational parameters, and observing forms. To order, send check or money order payable to "Timothy J. Robertson."
- **Lunar (Benton):** (1) *The ALPO Lunar Section's Selected Areas Program (SAP)*, \$17.50). Includes a full set of observing forms for the assigned or chosen lunar area or feature, together with a copy of the *Lunar Selected Areas Program Manual*. (2) *Observing Forms Packet*, \$10. Includes observing forms to replace those provided in the observing kit described above. Specify *Lunar Forms*. (See note for Venus.)
- **Mars (Troiani):** (1) *Martian Chronicle*; published approximately monthly during each apparition; send 8 to 10 SASEs; (2) *Observing Forms*; send SASE to obtain one form for you to copy; otherwise send \$3.60 to obtain 25 copies (make checks payable to "J.D. Beish.>").
- **Mars:** *ALPO Mars Observers Handbook*, send check or money order for \$9 per book (postage and handling included) to Astronomical League Sales, P.O. Box 572, West Burlington, IA 52655.
- **Jupiter:** (1) *Jupiter Observer's Startup Kit*, \$3 from the Jupiter Section Coordinator. (2) *Jupiter*, ALPO section newsletter, available online via the ALPO website or via snail-mail; send SASE's to the Jupiter Section Coordinator; (3) To join the ALPO Jupiter Section e-mail network, *J-Net*, send an e-mail message to the Jupiter Section Coordinator. (4) *Timing the Eclipses of Jupiter's Galilean Satellites* observ-

## ALPO Resources (continued)

- ing kit and report form; send SASE with 55 cents in postage stamps to John Westfall.
- **Saturn (Benton):** (1) *The ALPO Saturn Observing Kit*, \$20; includes introductory description of Saturn observing programs for beginners, a full set of observing forms, and a copy of *The Saturn Handbook*. (2) *Saturn Observing Forms Packet*, \$10; includes observing forms to replace those provided in the observing kit described above. Specify *Saturn Forms*. To order, see note for *Venus Forms*.
  - **Meteors:** (1) Pamphlet, *The ALPO Guide to Watching Meteors*, send check or money order for \$4 per book (postage and handling included) to Astronomical League Sales, P.O. Box 572, West Burlington, IA 52655. (2) *The ALPO Meteors Section Newsletter*, free (except postage), published quarterly (March, June, September, and December). Send check or money order for first class postage to cover desired number of issues to Robert D. Lunsford, 161 Vance St., Chula Vista, CA 91910.
  - **Minor Planets (Derald D. Nye):** *The Minor Planets Bulletin*, published quarterly \$9 per year in the U.S., Mexico and Canada, \$13 per year elsewhere (air mail only). Send check or money order payable to Mr. Nye to 10385 East Observatory Dr., Corona de Tucson, AZ 85641-2309.
  - **Computing Section (McClure):** Online newsletter, *The Digital Lens*, available via the World Wide Web and e-mail. To subscribe or make contributions, contact Mike McClure

## Other ALPO Publications

Checks must be in U.S. funds, payable to an American bank with bank routing number. Order from: ALPO, P.O. Box 2447, Antioch, CA 94531-2447 USA

- **An Introductory Bibliography for Solar System Observers.** Free for a stamped, self-addressed envelope. A 4-page list of books and magazines about Solar System bodies and how to observe them. The current edition was updated in October, 1998. Order from: ALPO Membership Secretary.
- **ALPO Membership Directory.** \$5 in North America; \$6 elsewhere. Latest updated list of members on 3.5-in. MS-DOS diskette; either DBASE or ASCII format. Make payment to "ALPO" Also available via e-mail as portable document format (pdf) file to requester's e-mail address. Provided at the discretion of the Membership Secretary. Order from Matthew Will, ALPO membership secretary/treasurer.
- **Back issues of *The Strolling Astronomer* (JALPO).** Many of the back issues listed below are almost out of stock, and it is impossible to guarantee that they will remain available. Issues will be sold on a first-come, first-served basis. In this list, volume

numbers are in italics, issue numbers are in plain type, and years are given in parentheses. The price is \$4 for each back issue; the current issue, the last one published, is \$5. We are always glad to be able to furnish old issues to interested persons and can arrange discounts on orders of more than \$30. Make payment to "Walter H. Haas" \$4 each: Vol. 8(1954); 7-8. Vol. 11 (1957); 11-12. Vol. 21(1968-69); 3-4 and 7-8. Vol. 23(1971-72); 7-8 and 9-10. Vol. 25(1974-76); 1-2, 3-4, and 11-12. Vol. 26 (1976.77); 3-4 and 11-12. Vol. 27(1977-79); 3-4 and 7-8. Vol. 31(1985-86); 9-10. Vol. 32(1987.88); 11.12. Vol. 33(1989); 7-9. Vol. 34(1990); 2 and 4. Vol. 37(1993-94); 1. Vol. 38 (1994-96); 1,2, and 3. Vol. 39(1996-97); 1, 3 and 4. Vol. 40(1998); 2. Vol. 41(1999); 4. Vol. 42 (2000); 1, 2, 3 and 4. Vol. 43 (2001); 1, 2 and current issue (43,3); \$5.00

## Membership in the ALPO

The **Association of Lunar and Planetary Observers (ALPO)** was founded by Walter H. Haas in 1947, and incorporated in 1990 as a medium for advancing and conducting astronomical work by both professional and amateur astronomers who share an interest in Solar-System observations. We welcome and provide services for all levels of astronomers: For the novice the **ALPO** is a place to learn and to enhance and practice techniques. For the advanced amateur it is a place where one's work will count. For the professional it is a resource where group studies or systematic observing patrols are necessary.

Our Association is an international group of students of the Sun, Moon, planets, asteroids, meteors, and comets. Our goals are to stimulate, coordinate, and generally promote the study of these bodies using methods and instruments that are available within the communities of both amateur and professional astronomers. We hold a conference each summer, usually in conjunction with other astronomical groups.

Membership dues include a subscription to the **Journal**. Memberships are for 1- or 2-year periods and include 4 or 8 consecutive issues, respectively.

The rates are \$23 US (1 year) or \$40 US (2 years) for the United States, Canada, and Mexico; and are \$30 US (1 year) or \$54 US (2 years) for all other countries. There is a 20-percent surcharge on all memberships obtained through subscription agencies or which require an invoice. Memberships for the online version of this Journal via the Internet are \$10 (1 year) or \$17 (US). Make payment by check or money order, payable (through a U.S. bank and encoded with U.S. standard banking numbers) to "**ALPO**"

# Mars:

**80 million miles away**

## Mars Globes: as close as your phone



### **Sky & Telescope's Mars Globe, 2nd Edition**

Created with more than 6,000 images taken by the Viking orbiters, our 12-inch globe approximates the planet's true color and emphasizes the striking bright and dark variations seen through telescopes.

**4676X ... \$99.95** plus shipping

### **Sky & Telescope's Mars Topography Globe**

This first-ever globe portrays the red planet's terrain in a color-coded elevation scheme that emphasizes mountains, valleys and craters. It incorporates more than 300 million measurements by the Mars Global Surveyor Spacecraft.

**46743 ... \$99.95** plus shipping

*Produced in cooperation with NASA and the U.S. Geological Survey, each of the 12-inch globes include 140 labeled features and is a scale model of the planet (1 inch = 350 miles or 570 kilometers).*

### **Special Offer:**

Save \$20 when you purchase both globes together! Place your order for the Mars Globe 2nd Edition and the Mars Topography Globe and pay \$179.90 (plus shipping.) Call 800-253-0245 or order securely online at [www.skypub.com](http://www.skypub.com).



**SKY**  
& TELESCOPE

**Call 800-253-0245**  
(Outside the U.S. and Canada  
+1 617-864-7360, fax: 617-864-6117  
[www.skypub.com](http://www.skypub.com))