

Meteor Activity Outlook for February 18-24, 2023




Julie Fearnley was photographing the southern sky from Kandanga, Queensland, Australia on August 20, 2022, at 19:45 AEST (9:45 UT) when this brilliant fireball shot through the Milky Way near the constellation of Crux (the Southern Cross). This is the same fireball that was posted last week, only this version has a much wider view. For more information on this event visit: https://ams.imo.net/members/imo_view/event/2022/5347 ©Julie Fearnley


During this period, the moon reaches its new phase on Monday February 20th. At that time the moon will lie near the sun and will be invisible at night. As the week progresses the moon will enter the evening sky with a thin crescent phase. The thin phase and the fact that the moon will set long before the more active morning hours arrive means that this is the best time to view meteor activity during February. The estimated total hourly rates for evening observers this week should be near 3 as seen from mid-northern latitudes (45N) and 4 as seen from tropical southern locations (25S) For morning observers, the estimated total hourly rates should be near 9 as seen from mid-northern latitudes (45N) and 16 as seen from tropical southern locations (25S). The actual rates seen will also depend on factors such as personal light and motion perception, local weather conditions, alertness, and experience in watching meteor activity. Note that the hourly rates listed below are estimates as viewed from dark sky sites away from urban light sources. Observers viewing from urban areas will see less activity as only the brighter meteors will be visible from such locations.

The radiant (the area of the sky where meteors appear to shoot from) positions and rates listed below are exact for Saturday night/Sunday morning February 18/19. These positions do not change greatly day to day so the listed coordinates may be used during this entire period. Most star atlases (available at science stores and planetariums) will provide maps with grid lines of the celestial coordinates so that you may find out exactly where these positions are located in the sky. I have also included charts of the sky that display the radiant positions for evening, midnight, and morning. The center of each chart is the sky directly overhead at the appropriate hour. These charts are oriented for facing south but can be used for any direction by rotating the charts to the


desired direction. A planisphere or computer planetarium program is also useful in showing the sky at any time of night on any date of the year. Activity from each radiant is best seen when it is positioned highest in the sky, either due north or south along the meridian, depending on your latitude. It must be remembered that meteor activity is rarely seen at the radiant position. Rather they shoot outwards from the radiant, so it is best to center your field of view so that the radiant lies at the edge and not the center. Viewing there will allow you to easily trace the path of each meteor back to the radiant (if it is a shower member) or in another direction if it is sporadic. Meteor activity is not seen from radiants that are located far below the horizon. The positions below are listed in a west to east manner in order of right ascension (celestial longitude). The positions listed first are located further west therefore are accessible earlier in the night while those listed further down the list rise later in the night.



Radiant Positions at 8pm Local Standard Time



Radiant Positions at Midnight Local Standard Time



Radiant Positions at 4am Local Standard Time

These sources of meteoric activity are expected to be active this week.

The large **Anthelion (ANT)** is currently centered at 10:52 (163) +07. This position lies in southern Leo, 10 degrees southeast of the 1st magnitude star known as Regulus (alpha Leonis). Due to the large size of this radiant, these meteors may also be seen from northeastern Sextans as well as Leo. This radiant is best placed near 01:00 local standard time (LST) when it lies on the meridian and is highest in the southern sky. Rates at this time should be near 3 per hour no matter your location. With an entry velocity of 30 km/sec., the average Anthelion meteor would be of slow velocity.

The last of the **alpha Centaurids (ACE)** should appear this weekend. The radiant is currently located at 14:56 (224) -62. This position lies in the faint constellation of Circinus, 2 degrees southeast of the zero-magnitude star known as Rigel Kentaurus (alpha Centauri). Due to the southern declination of this radiant, these meteors are not well seen in the northern hemisphere. Current hourly rates are expected to be less than 1 no matter your location. These meteors are best seen near 05:00 LST when the radiant lies highest above the southern horizon. At 56 km/sec. the alpha Centaurids would produce mostly swift meteors.

As seen from the mid-northern hemisphere (45N) one would expect to see approximately 6 **sporadic** meteors per hour during the last hour before dawn as seen from rural observing sites. Evening rates would be near 2 per hour. As seen from the tropical southern latitudes (25S), morning rates would be near 14 per hour as seen from rural observing sites and 4 per hour during the evening hours.

You can keep track of the activity of these meteor showers as well as those beyond the limits of visual observing by visiting the [NASA Meteor Shower Portal](#). You can move the sky globe to see different areas of the sky. Colored dots indicate shower meteors while white dots indicate sporadic (random) activity. The large orange disk indicates the position of the sun so little activity will be seen in that area of the sky.

The list below offers the information in tabular form of the showers that I feel are within reach of the visual observer to discern. Hourly rates are often less than 1 but noting parameters such as the radiant distance and the elevation of each meteor, one can compute the probability of shower association. Most showers discovered by video means have rates less than 1 per **night** away from maximum, so the showers listed in these outlooks are not as weak as they seem. Rates and positions are exact for Saturday night/Sunday morning except where noted in the shower descriptions.

SHOWER	DATE OF MAXIMUM ACTIVITY	CELESTIAL POSITION	ENTRY VELOCITY	CULMINATION	HOURLY RATE	CLASS
		RA (RA in Deg.) DEC	Km/Sec	Local Standard Time	North- South	
Anthelions (ANT)	-	10:52 (163) +07	30	01:00	3 - 3	II
alpha Centaurids (ACE)	Feb 08	14:56 (224) -62	59	05:00	<1 - <1	II

Class Explanation: A scale to group meteor showers by their intensity:

- **Class I:** the strongest annual showers with Zenith Hourly Rates normally ten or better.
- **Class II:** reliable minor showers with ZHR's normally two to ten.
- **Class III:** showers that do not provide annual activity. These showers are rarely active yet have the potential to produce a major display on occasion.
- **Class IV:** weak minor showers with ZHR's rarely exceeding two. The study of these showers is best left to experienced observers who use plotting and angular velocity estimates to determine shower association. These weak showers are also good targets for video and photographic work. Observers with less experience are urged to limit their shower associations to showers with a rating of I to III.