

# Meteor Activity Outlook for 25-31 March 2023




Bart Benjamin captured this bright fireball just above his house on August 2, 2022, at 22:34 CDT (3:34 UT on August 3rd) from Bloomingdale, Illinois, USA. Bart mentions that his image was taken with an Olympus OM-D E-M5 Mark III camera with Olympus M.Zuiko 12 mm lens, mounted on a tripod in my backyard facing north toward my house. Olympus' live composite feature slowly build a composite image (in camera) of 2.5 second exposures, only registering changes to each pixel brightness, building a nice star trail image without overexposing the overall frame. A total of 1,441 frames were taken, spanning 60 minutes from 22:14 to 23:14 CDT. For more information on this event visit: [https://ams.imo.net/members/imo\\_view/event/2022/4637](https://ams.imo.net/members/imo_view/event/2022/4637) ©Bart Benjamin

During this period, the moon reaches its first quarter phase on Wednesday March 29th. At that time the moon will lie 90 degrees east of the sun and will set near 0300 local daylight saving time (LDST). Observing may be slightly compromised by the waxing moon, but once the moon sets viewing circumstances are optimal. The estimated total hourly rates for evening observers this week should be near 2 as seen from mid-northern latitudes (45N) and 3 as seen from tropical southern locations (25S) For morning observers, the estimated total hourly rates should be near 7 as seen from mid-northern latitudes (45N) and 11 as seen from tropical southern locations (25S). The actual rates seen will also depend on factors such as personal light and motion perception, local weather conditions, alertness, and experience in watching meteor activity. Evening and early morning rates are slightly reduced due to moonlight. Note that the hourly rates listed below are estimates as viewed from dark sky sites away from urban light sources. Observers viewing from urban areas will see less activity as only the brighter meteors will be visible from such locations.

The radiant (the area of the sky where meteors appear to shoot from) positions and rates listed below are exact for Saturday night/Sunday morning March 25/26. These positions do not change greatly day to day so the listed


coordinates may be used during this entire period. Most star atlases (available at science stores and planetariums) will provide maps with grid lines of the celestial coordinates so that you may find out exactly where these positions are located in the sky. I have also included charts of the sky that display the radiant positions for evening, midnight, and morning. The center of each chart is the sky directly overhead at the appropriate hour. These charts are oriented for facing south but can be used for any direction by rotating the charts to the desired direction. A planisphere or computer planetarium program is also useful in showing the sky at any time of night on any date of the year. Activity from each radiant is best seen when it is positioned highest in the sky, either due north or south along the meridian, depending on your latitude. It must be remembered that meteor activity is rarely seen at the radiant position. Rather they shoot outwards from the radiant, so it is best to center your field of view so that the radiant lies at the edge and not the center. Viewing there will allow you to easily trace the path of each meteor back to the radiant (if it is a shower member) or in another direction if it is sporadic. Meteor activity is not seen from radiants that are located far below the horizon. The positions below are listed in a west to east manner in order of right ascension (celestial longitude). The positions listed first are located further west therefore are accessible earlier in the night while those listed further down the list rise later in the night.



## Radiant Positions at 9pm Local Daylight Saving Time



Radiant Positions at 1am Local Daylight Saving Time



Radiant Positions at 5am Local Daylight Saving Time

## **These sources of meteoric activity are expected to be active this week.**

The large **Anthelion (ANT)** radiant is currently centered at 13:12 (198) -07. This position lies in southern Virgo, 4 degrees northwest of the 1st magnitude star known as Spica (alpha Virginis). Due to the large size of this radiant, these meteors may also be seen from northern Corvus as well as Virgo. This radiant is best placed near 02:00 LDST when it lies on the meridian and is highest in the southern sky. Rates at this time should be near 2 per hour no matter your location. With an entry velocity of 30 km/sec., the average Anthelion meteor would be of slow velocity.

The **gamma Normids (GNO)** are an ill-defined source, only well seen from the Southern Hemisphere. Some sources dispute its existence while others have differing activity periods in March. Here we list the parameters from an article on Meteor showers from the Southern Hemisphere by Sirko Molau and Steve Kerr\*. Their results show the activity period is from March 22-27 with no distinct maximum. On March 25th the radiant is located at 15:56 (239) -51. This area of the sky lies in central Norma, 4 degrees west of the 4<sup>th</sup> magnitude star known as Gamma<sup>2</sup> Normae. These meteors are best seen during the last dark hour before dawn, when the radiant lies highest in a dark sky. With an entry velocity of 56km/sec., the average meteor from this source would be of fast velocity. Expected rates are less than 1 per hour as seen from the Northern Hemisphere and near 1 as seen from south of the equator. These meteors are poorly seen from the Northern Hemisphere.

\*Molau, Sirko and Kerr, Steve, WGN 42-2 (2014) Meteor Showers of the Southern Hemisphere, page 25

The **delta Pavonids (DPA)** were discovered by Michael Buhagiar from Australia in the 1970's. These meteors are active from March 11 through April 16, with maximum activity occurring on March 31st. The current position of the radiant lies near 20:00 (300) -62. This area of the sky lies in northern Pavo, 6 degrees southwest of the 2<sup>nd</sup> magnitude star known as Peacock (alpha Pavonis). These meteors are best seen during the last dark hour prior to dawn when the radiant lies in the southeastern sky. With an entry velocity of 58km/sec., the average meteor from this source would be of fast velocity. Expected rates are less than 1 per hour during this period. Like the gamma Normids, these meteors are poorly seen from the Northern Hemisphere.

As seen from the mid-northern hemisphere (45N) one would expect to see approximately 5 **sporadic** meteors per hour during the last hour before dawn as seen from rural observing sites. Evening rates would be near 1 per hour. As seen from the tropical southern latitudes (25S), morning rates would be near 8 per hour as seen from rural observing sites and 2 per hour during the evening hours. Evening rates are slightly reduced by moonlight during this period.

You can keep track of the activity of these meteor showers as well as those beyond the limits of visual observing by visiting the [NASA Meteor Shower Portal](#). You can move the sky globe to see different areas of the sky. Colored dots indicate shower meteors while white dots indicate sporadic (random) activity. The large orange disk indicates the position of the sun so little activity will be seen in that area of the sky.

The list below offers the information in tabular form of the showers that I feel are within reach of the visual observer to discern. Hourly rates are often less than 1 but noting parameters such as the radiant distance and the elevation of each meteor, one can compute the probability of shower association. Most showers discovered by video means have rates less than 1 per **night** away from maximum, so the showers listed in these outlooks are not as weak as they seem. Rates and positions are exact for Saturday night/Sunday morning except where noted in the shower descriptions.

SHOWER	DATE OF MAXIMUM ACTIVITY	CELESTIAL POSITION	ENTRY VELOCITY	CULMINATION	HOURLY RATE	CLASS
		RA (RA in Deg.) DEC	Km/Sec	Local Daylight Saving Time	North- South	
Anthelions (ANT)	-	13:12 (198) -07	30	02:00	2 - 2	II
gamma Normids (GNO)	Mar 25	15:56 (239) -51	56	05:00	<1 - 1	IV
delta Pavonids (DPA)	Mar 31	20:00 (300) -62	58	09:00	<1 - <1	IV

**Class Explanation:** A scale to group meteor showers by their intensity:

- **Class I:** the strongest annual showers with Zenith Hourly Rates normally ten or better.
- **Class II:** reliable minor showers with ZHR's normally two to ten.
- **Class III:** showers that do not provide annual activity. These showers are rarely active yet have the potential to produce a major display on occasion.
- **Class IV:** weak minor showers with ZHR's rarely exceeding two. The study of these showers is best left to experienced observers who use plotting and angular velocity estimates to determine shower association. These weak showers are also good targets for video and photographic work. Observers with less experience are urged to limit their shower associations to showers with a rating of I to III.