

Meteor Activity Outlook for June 17-23, 2023




Ryan Conner captured this multi-bursting Southern Taurid fireball using his [AllSky Camera System](#) on October 29, 2022, at 19:53 EDT (23:53 UT) from North Royalton, Ohio, USA. The five dots in a line crossing the fireball belong to an aircraft. ©Ryan Conner

During this period, the moon reaches its new phase on Sunday June 18th. At that time, it will lie near the sun and will be invisible at night. As the week progresses the thin crescent moon will enter the evening sky but will set during the evening hours and will not interfere with meteor observing during the more active morning hours. The estimated total hourly rates for evening observers this week should be near 3 as seen from mid-northern latitudes (45N) and 4 as seen from tropical southern locations (25S) For morning observers, the estimated total hourly rates should be near 10 as seen from mid-northern latitudes (45N) and 12 as seen from tropical southern locations (25S). The actual rates seen will also depend on factors such as personal light and motion perception, local weather conditions, alertness, and experience in watching meteor activity. Note that the hourly rates listed below are estimates as viewed from dark sky sites away from urban light sources. Observers viewing from urban areas will see less activity as only the brighter meteors will be visible from such locations.

The radiant (the area of the sky where meteors appear to shoot from) positions and rates listed below are exact for Saturday night/Sunday morning June 17/18. These positions do not change greatly day to day so the listed coordinates may be used during this entire period. Most star atlases (available at science stores and planetariums) will provide maps with grid lines of the celestial coordinates so that you may find out exactly where these positions are located in the sky. I have also included charts of the sky that display the radiant positions for evening, midnight, and morning. The center of each chart is the sky directly overhead at the appropriate hour. These charts are oriented for facing south but can be used for any direction by rotating the charts to the desired direction. A planisphere or computer planetarium program is also useful in showing the sky at any time of night on any date of the year. Activity from each radiant is best seen when it is positioned highest in the sky, either due north or south along the meridian, depending on your latitude. It must be remembered that meteor activity is rarely seen at its radiant position. Rather

they shoot outwards from the radiant, so it is best to center your field of view so that the radiant lies at the edge and not the center. Viewing there will allow you to easily trace the path of each meteor back to the radiant (if it is a shower member) or in another direction if it is sporadic. Meteor activity is not seen from radiants that are located far below the horizon. The positions below are listed in a west to east manner in order of right ascension (celestial longitude). The positions listed first are located further west therefore are accessible earlier in the night while those listed further down the list rise later in the night.



Radiant Positions at 10pm Local Daylight-Saving Time



Radiant Positions at 1am Local Daylight-Saving Time



Radiant Positions at 4am Local Daylight-Saving Time

These sources of meteoric activity are expected to be active this week.

The center of the large **Anthelion (ANT)** radiant is currently located at 18:36 (279) -23. This position lies in northern Sagittarius, 3 degrees northeast of the 3rd magnitude star known as Kaus Borealis (λ Sagittarii). Due to the large size of this radiant, Anthelion activity may also appear from southern Scutum as well as Sagittarius. This radiant is best placed near 0100 LST, when it lies on the meridian and is located highest in the sky. Rates at this time should be near 2 per hour as seen from the northern hemisphere and 3 per hour as seen from south of the equator. With an entry velocity of 30 km/sec., the average Anthelion meteor would be of slow velocity.

The **phi Piscids (PPS)** have been found to be comprised of two components with separate activity periods. Component A is active from June 13 through July 5 with maximum activity occurring on June 25th. Current rates are expected to be near 1 meteor per hour just before dawn. The position of the radiant lies near 00:19 (005) +18. This position lies in northern Pisces, 4 degrees northeast of the 3rd magnitude star known as Algenib (γ Pegasi). This area of the sky lies highest in the northeastern sky during the last hour prior to dawn. With an entry velocity of 66km/sec., these meteors are swift.

The **c-Andromedids (CAN)** were discovered by Sirko Molau and Juergen Rendtel using video data from the IMO network. Activity from this source is seen from June 21 through July 28 with maximum activity occurring on July 12. On June 21st the radiant will lie at 00:43 (011) +41, which places it in central Andromeda, within the Great Andromeda Galaxy, which is visible to the naked eye as a small oval cloud. This area of the sky is best seen during the last dark hour before dawn when the radiant lies highest in the northern sky. Observers in the northern hemisphere are better situated to view this activity as the radiant rises much higher in the sky before dawn compared to southern latitudes. Current rates would be less than 1 per hour no matter your location. With an entry velocity of 56 km/sec., the average meteor from this source would be of swift velocity.

As seen from the mid-northern hemisphere (45N) one would expect to see approximately 7 **sporadic** meteors per hour during the last hour before dawn as seen from rural observing sites. Evening rates would be near 2 per hour. As seen from the tropical southern latitudes (25S), morning rates would be near 9 per hour as seen from rural observing sites and 3 per hour during the evening hours. Locations between these two extremes would see activity between the listed figures.

You can keep track of the activity of these meteor showers as well as those beyond the limits of visual observing by visiting the [NASA Meteor Shower Portal](#). You can move the sky globe to see different areas of the sky. Colored dots indicate shower meteors while white dots indicate sporadic (random) activity. The large orange disk indicates the position of the sun so little activity will be seen in that area of the sky.

The list below offers the information in tabular form of the showers that I feel are within reach of the visual observer to discern. Hourly rates are often less than 1 but noting parameters such as the radiant distance and the elevation of each meteor, one can compute the probability of shower association. Most showers discovered by video means have rates less than 1 meteor per **night** away from maximum, so the showers listed in these articles are not as weak as they seem. Rates and

positions are exact for Saturday night/Sunday morning except where noted in the shower descriptions.

SHOWER	DATE OF MAXIMUM ACTIVITY	CELESTIAL POSITION	ENTRY VELOCITY	CULMINATION	HOURLY RATE	CLASS
Anthelion (ANT)	-	18:36 (279) -23	30	02:00	2 - 3	II
phi Piscids (PPS)	Jun 25	00:19 (005) +18	66	08:00	1 - 1	IV
c- Andromedids (CAN)	Jul 12	00:43 (011) +41	56	09:00	<1 - <1	IV

Class Explanation: A scale to group meteor showers by their intensity:

- **Class I:** the strongest annual showers with Zenith Hourly Rates normally ten or better.
- **Class II:** reliable minor showers with ZHR's normally two to ten.
- **Class III:** showers that do not provide annual activity. These showers are rarely active yet have the potential to produce a major display on occasion.
- **Class IV:** weak minor showers with ZHR's rarely exceeding two. The study of these showers is best left to experienced observers who use plotting and angular velocity estimates to determine shower association. These weak showers are also good targets for video and photographic work. Observers with less experience are urged to limit their shower associations to showers with a rating of I to III.